UT-Komaba/12-9
TIT/HEP-625

arXiv:1211.1624v2 [hep-th] 2 Dec 2012

Effective field theory approach to quasi-single field inflation

Toshifumi Noumi,[1, *] Masahide Yamaguchi,[2, †] and Daisuke Yokoyama[2, ‡]

[1] *Institute of Physics, University of Tokyo, Komaba, Meguro-ku, Tokyo 153-8902, Japan*
[2] *Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan*

We apply the effective field theory approach to quasi-single field inflation, which contains an additional scalar field with Hubble scale mass other than inflaton. Based on the time-dependent spatial diffeomorphism, which is not broken by the time-dependent background evolution, the most generic action of quasi-single field inflation is constructed up to third order fluctuations. Using the obtained action, the effects of the additional massive scalar field on the primordial curvature perturbations are discussed. In particular, we calculate the power spectrum and discuss the momentum-dependence of three point functions in the squeezed limit for general settings of quasi-single field inflation. Our framework can be also applied to inflation models with heavy particles. We make a qualitative discussion on the effects of heavy particles during inflation and that of sharp turning trajectory in our framework.

I. INTRODUCTION

Inflation gives the most natural solution to the horizon and the flatness problems of the big-bang theory as well as generates the primordial perturbations [1, 2], whose properties well coincide with the recent observations of cosmic microwave background anisotropies like the Wilkinson Microwave Anisotropy Probe [3]. Models of inflation can be classified into two categories with respect to relevant degrees of freedom during inflation: single-field models and multiple field models. Recently, the most general single field inflation model with the second order equations of motion [4] has been invented in the context of Horndeski [5, 6] and Galileon theories [7, 8]. Then, the bispectra of primordial curvature [9, 10] and tensor perturbations [11] are obtained as well as their powerspectra [4].

Effective scalar fields are ubiquitous in the extensions of the standard model of particle physics such as supergravity and superstring. Then, it is well motivated to consider multiple field models of inflation. Such multiple field models are roughly divided into three classes: (i) only one field is light, while the other fields are very heavy compared to the Hubble scale during inflation. Generically, this class virtually falls into the single field category [12]. However, it is recently discussed that heavy modes can affect the dynamics of light mode in some particular cases [13–20]. (ii) there are multiple light fields, in which isocurvature perturbations are generated in addition to curvature perturbations. (iii) only one field is light, while the masses of other fields are comparable to the Hubble scale during inflation. This class is called quasi-single field inflation model [21, 22].

In supergravity, inflation necessarily involves supersymmetry (SUSY) breaking, whose effects are transmitted into other scalar fields as Hubble induced masses [23–25].[1] Therefore, quasi-single field inflation is naturally realized in supergravity and it is well motivated by the model building based on supergravity or inspired by superstring. Furthermore, it is known that massive isocurvature modes which couple to the inflaton and have Hubble scale masses can give significant impacts on primordial curvature perturbations. In the original paper [21] by Chen and Wang, it was shown that, for example, scalar three point functions take the intermediate shapes between local and equilateral types. Based on these backgrounds, we would like to discuss quasi-single field inflation model in general settings.

Recently, effective field theory approach to inflation has been invented in [27–29], which is based on the symmetry breaking during inflation: Time diffeomorphism is broken by the time-dependent background evolution during inflation. Then, based on the unbroken time-dependent spatial diffeomorphism, the effective action for inflation can be constructed systematically in unitary gauge, where inflaton is eaten by graviton and there are no perturbations of inflaton. By use of the Stückelberg trick, the curvature perturbation can be associated with the Goldstone boson π, which non-linearly realizes the time diffeomorphism. The key observation is that the Goldstone π could decouple from the metric fluctuations in some parameter region which we call the decoupling regime. In the decoupling regime,

---

[*] Email: tnoumi"at"hep1.c.u-tokyo.ac.jp
[†] Email: gucci"at"phys.titech.ac.jp
[‡] Email: d.yokoyama"at"th.phys.titech.ac.jp

[1] The methods to keep inflaton flat against such SUSY breaking effects are reviewed in Refs. [26].

the dynamics of Goldstone π is described by a simplified action, which does not contain metric perturbations. As a consequence, calculations of scalar perturbations are also simplified and seeds of non-Gaussianities become clear.

In this paper, we apply this effective field theory approach to quasi-single field inflation. First, in unitary gauge, we write down the most general action invariant under the time-dependent spatial diffeomorphism and constructed from graviton and the massive isocurvature mode. The obtained action is expanded systematically in fluctuations and derivatives around the FRW background. By the Stückelberg trick, we introduce the the action for Goldstone boson and carefully discuss its decoupling regime. Using the action in the decoupling regime, the power spectrum is calculated in the general setting of quasi-single field inflation. The momentum dependence of scalar three-point function is also discussed in the general setting. Our framework can be also applied to inflation models with heavy particles. As an application, we make a qualitative discussion on the effects of heavy particles during inflation and that of sharp turning trajectory.

The organization of this paper is as follows. In the next section, we briefly review the effective field theory approach and quasi-single field inflation. In Sec. III, the most general action for quasi-single field inflation is constructed via effective field theory approach. The decoupling regime of the obtained action is also discussed. In Sec. IV, the power spectrum is calculated first in the general setting of quasi-single field inflation with constant mixing couplings. Then, the effects of sharp turning trajectory on the power spectrum is qualitatively discussed. In Sec. V, the momentum dependence of scalar three-point functions are discussed. Final section is devoted to summary and discussions. Technical details of the calculation of the power spectrum are summarized in Appendices.

II. EFFECTIVE FIELD THEORY APPROACH AND QUASI-SINGLE FIELD INFLATION

In this section we briefly review the effective field theory approach to inflation developed in [27] and the quasi-single field inflation model proposed in [21].

A. Effective field theory approach to inflation

Inflation is an accelerated cosmic expansion with an approximately constant Hubble parameter:

$$ds^2 = -dt^2 + a^2(t)dx^i dx^i \quad \text{with} \quad H(t) = \frac{\dot{a}}{a}\,, \quad \epsilon = -\frac{\dot{H}}{H^2} \ll 1\,. \tag{1}$$

It is characterized by the spontaneous breaking of the time-diffeomorphism:

$$\langle \phi(t,x) \rangle = \phi_0(t)\,, \tag{2}$$

where $\phi(t,x)$ is some scalar operator. Here we chose the frame in which the vacuum expectation value of $\phi(t,x)$ is spatially uniform. Assuming the degrees of freedom relevant to the cosmological perturbation and invariance under the time-dependent spatial diffeomorphism, $x^i \to x'^{\,i} = x^i + \xi^i(t,x^j)$, which is not broken by the condensation $\phi_0(t)$, we can construct the effective action for inflation.

In the simplest case, relevant degrees of freedom are three physical modes of graviton: two transverse modes and one longitudinal mode related to the inflaton. As discussed in [27], any action of graviton invariant under the time-dependent spatial diffeomorphism can be written in terms of the Riemann tensor $R_{\mu\nu\rho\sigma}$, the time-like component of the metric g^{00}, the extrinsic curvature $K_{\mu\nu}$ on constant-t surfaces, the covariant derivative ∇_μ, and the time coordinate t:

$$S = \int d^4x\sqrt{-g}\,F(R_{\mu\nu\rho\sigma}, g^{00}, K_{\mu\nu}, \nabla_\mu, t)\,, \tag{3}$$

where all the free indices inside the function F must be upper 0's. Note that g^{00} should be treated as a scalar when considering its covariant derivative, and we can use $\partial_\mu g^{00}$ for example. The explicit form of the extrinsic curvature $K_{\mu\nu}$ is

$$K_{\mu\nu} = h_\mu^\sigma \nabla_\sigma n_\nu = -\frac{\delta_\mu^0 \partial_\nu g^{00} + \delta_\nu^0 \partial_\mu g^{00}}{2(-g^{00})^{3/2}} - \frac{\delta_\mu^0 \delta_\nu^0 g^{0\sigma}\partial_\sigma g^{00}}{2(-g^{00})^{5/2}} + \frac{g^{0\rho}(\partial_\mu g_{\rho\nu} + \partial_\nu g_{\rho\mu} - \partial_\rho g_{\mu\nu})}{2(-g^{00})^{1/2}}\,, \tag{4}$$

where $n_\mu = -\dfrac{\delta^0_\mu}{\sqrt{-g^{00}}}$ is a unit vector perpendicular to constant t surfaces and $h_{\mu\nu} = g_{\mu\nu} + n_\mu n_\nu$ is the induced spatial metric on constant t surfaces. In [27], it was shown that the action (3) can be expanded around a given FRW background as

$$S = \int d^4x\sqrt{-g}\left[\frac{1}{2}M_{\rm Pl}^2 R + M_{\rm Pl}^2\dot{H}g^{00} - M_{\rm Pl}^2(3H^2+\dot{H}) + F^{(2)}(\delta g^{00}, \delta K_{\mu\nu}, \delta R_{\mu\nu\rho\sigma}; \delta^0_\mu, g_{\mu\nu}, g^{\mu\nu}, \nabla_\mu, t)\right], \tag{5}$$

where the function $F^{(2)}$ starts with quadratic terms of the arguments δg^{00}, $\delta K_{\mu\nu}$, and $\delta R_{\mu\nu\rho\sigma}$ and all the free indices must be upper 0's. The arguments δg^{00}, $\delta K_{\mu\nu}$, and $\delta R_{\mu\nu\rho\sigma}$ are defined by[2]

$$\delta g^{00} = g^{00} + 1\,, \tag{6}$$

$$\delta K_{\mu\nu} = K_{\mu\nu} - H h_{\mu\nu}\,, \tag{7}$$

$$\delta R_{\mu\nu\rho\sigma} = R_{\mu\nu\rho\sigma} - 2H^2 h_{\mu[\rho}h_{\sigma]\nu} + (\dot{H}+H^2)(h_{\mu\rho}\delta^0_\nu\delta^0_\sigma + (3\text{ permutations}))\,. \tag{8}$$

They are covariant under time-dependent spatial-diffeomorphism and vanish on the FRW background. Notice that the action of single field inflation in the uniform inflaton gauge can be reproduced by gauge-fixing the time-dependent spatial diffeomorphism as

$$g_{ij}(x) = a^2(t)e^{2\zeta(x)}(e^{\gamma(x)})_{ij} \quad \text{with} \quad \gamma_{ii} = \partial_i\gamma_{ij} = 0\,, \tag{9}$$

where $\zeta(x)$ is the scalar perturbation.

For the calculation of correlation functions of the scalar perturbation ζ, it is convenient to introduce the action for the Goldstone boson π by the Stückelberg method. We perform the following time-diffeomorphism on the action (5) in the unitary gauge:

$$t \to \tilde{t}\,, \quad x^i \to \tilde{x}^i \quad \text{with} \quad \tilde{t} + \tilde{\pi}(\tilde{t},\tilde{x}) = t\,, \quad \tilde{x}^i = x^i\,. \tag{10}$$

In general, the transformation (10) is realized by the following replacement:

$$\begin{aligned}&\delta^0_\mu \to \delta^0_\mu + \partial_\mu\pi\,, \quad f(t) \to f(t+\pi)\,, \quad \int d^4x\sqrt{-g} \to \int d^4x\sqrt{-g}\,,\\ &\nabla_\mu \to \nabla_\mu\,, \quad g_{\mu\nu} \to g_{\mu\nu}\,, \quad g^{\mu\nu} \to g^{\mu\nu}\,, \quad R_{\mu\nu\rho\sigma} \to R_{\mu\nu\rho\sigma}\,,\end{aligned} \tag{11}$$

where we dropped the tilde for simplicity and g^{00} transforms, for example, as

$$g^{00} \to g^{00} + 2g^{0\mu}\partial_\mu\pi + g^{\mu\nu}\partial_\mu\pi\partial_\nu\pi\,. \tag{12}$$

The transformation rules of $K_{\mu\nu}$ and $h_{\mu\nu}$ also follow from (11) straightforwardly. These procedures lead to the following action for the Goldstone boson π:

$$S = \int d^4x\sqrt{-g}\left[\frac{1}{2}M_{\rm Pl}^2 R + M_{\rm Pl}^2\dot{H}(t+\pi)\left(g^{00} + 2g^{0\mu}\partial_\mu\pi + g^{\mu\nu}\partial_\mu\pi\partial_\nu\pi\right) - M_{\rm Pl}^2\Big(3H^2(t+\pi) + \dot{H}(t+\pi)\Big) + \ldots\right], \tag{13}$$

where the dots stand for the terms corresponding to $F^{(2)}$. The obtained action enjoys the time-diffeomorphism by assigning to π the non-linear transformation rule[3]

$$\pi(x) \to \tilde{\pi}(\tilde{x}) = \pi(x) - \xi^0(x) \quad \text{with} \quad t \to \tilde{t} = t + \xi^0(x)\,, \quad x^i \to \tilde{x}^i = x^i\,, \tag{14}$$

and the action in the unitary gauge can be reproduced by gauge-fixing the time-diffeomorphism as $\pi(x) = 0$.

---

[2] Here and in what follows, we concentrate on the spatially flat FRW background.

[3] Here it should be noted that while the action is invariant under the time-diffeomorphism, it does not have the shift symmetry $\pi \to \pi + \text{constant}$ because of the time-dependent free parameters such as $H(t+\pi)$ or $\dot{H}(t+\pi)$. Expanding the parameters in π, we find, for example, that π has a mass term $\sim M_{\rm Pl}^2\dot{H}^2\pi^2$, which is sub-leading in the slow-roll approximation.

It is important to recognize that, in the action (13), terms with graviton fluctuations have less derivatives than those without graviton. Because of this property, it is expected that the mixing of the Goldstone boson and graviton becomes irrelevant to the dynamics of the Goldstone boson at a sufficiently high energy scale. For example, let us consider the following simplest case:

$$S=\int d^4x\sqrt{-g}\left[\frac{1}{2}M_{\rm Pl}^2R+M_{\rm Pl}^2\dot{H}(t+\pi)\left(g^{00}+2g^{0\mu}\partial_\mu\pi+g^{\mu\nu}\partial_\mu\pi\partial_\nu\pi\right)-M_{\rm Pl}^2\Big(3H^2(t+\pi)+\dot{H}(t+\pi)\Big)\right]\,. \tag{15}$$

In the canonical normalization ($\pi_c\sim M_{\rm Pl}(-\dot{H})^{1/2}\pi$, $\delta g_c^{\mu\nu}\sim M_{\rm Pl}\delta g^{\mu\nu}$), the mixing term $M_{\rm Pl}^2\,\dot{H}\,\delta g^{0\mu}\,\partial_\mu\pi$ can be written as

$$M_{\rm Pl}^2\,\dot{H}\,\delta g^{0\mu}\,\partial_\mu\pi\sim(-\dot{H})^{1/2}\,\delta g_c^{0\mu}\,\partial_\mu\pi_c\sim\epsilon^{1/2}H\,\delta g_c^{0\mu}\,\partial_\mu\pi_c\,, \tag{16}$$

and it can be neglected in the energy scale $E\gg\epsilon^{1/2}H$. In other words, when the slow-roll parameter ϵ is small, the mixing becomes irrelevant inside the horizon. The dynamics of π inside the horizon are then determined by the following action in the decoupling limit:

$$S\sim\int d^4x\,a^3\Big[M_{\rm Pl}^2\dot{H}(t+\pi)\Big(-1-2\dot{\pi}-\dot{\pi}^2+\frac{(\partial_i\pi)^2}{a^2}\Big)-M_{\rm Pl}^2(3H^2(t+\pi)+\dot{H}(t+\pi))\Big]\,, \tag{17}$$

where the metric reduces to the FRW background. More generally, the mixing of the Goldstone boson and graviton becomes irrelevant inside the horizon when the free parameters of the action are in some regime (decoupling regime) as in the slow-roll regime for the above simplest case. In the decoupling regime, the calculation of scalar correlation functions also becomes tractable. Taking the spatially flat gauge[4]

$$g_{ij}(x)=a^2(t)(e^{\gamma(x)})_{ij}\quad\text{with}\quad\gamma_{ii}=\partial_i\gamma_{ij}=0\,, \tag{18}$$

the scalar perturbation $\zeta(x)$ is given by $\zeta(x)=-H(t)\pi(x)$ at the linear order, and the calculation of correlation functions of ζ reduces to those of π, which can be obtained using the simplified action in the decoupling limit. This kind of simplification in the decoupling regime is one of the advantages to use the effective field theory approach. In the next section we extend this approach to quasi-single field inflation.

B. Quasi-single field inflation

The original model [21] of quasi-single field inflation is described by the following matter action:

$$S_{\rm matter}=\int d^4x\sqrt{-g}\Big[-\frac{1}{2}(\tilde{R}+\chi)^2g^{\mu\nu}\partial_\mu\theta\partial_\nu\theta-\frac{1}{2}g^{\mu\nu}\partial_\mu\chi\partial_\nu\chi-V_{\rm sr}(\theta)-V(\chi)\Big]\,, \tag{19}$$

where θ and χ are the tangential and radial directions of a circle with radius $\tilde{R}$ and the potential $V_{\rm sr}(\theta)$ along the tangential direction is of slow-roll type. The homogeneous backgrounds and their equations of motion are given by

$$\begin{aligned}&\theta=\theta_0(t)\,,\quad\chi=\chi_0\,(\text{constant})\,,\\&3M_{\rm Pl}^2H^2=\frac{1}{2}R^2\theta_0^2+V(\chi_0)+V_{\rm sr}(\theta_0)\,,\quad-2M_{\rm Pl}^2\dot{H}=R^2\dot{\theta}_0^2\,,\quad V'(\chi_0)=R\dot{\theta}_0^2\,,\quad R^2\ddot{\theta}_0+3R^2H\dot{\theta}_0+V_{\rm sr}'(\theta_0)=0\,,\end{aligned} \tag{20}$$

where $R=\tilde{R}+\chi_0$. Expanding the action around the homogeneous background, it yields the following second order action of the fluctuations $\delta\theta=\theta-\theta_0$ and $\sigma=\chi-\chi_0$:

$$S_{\rm matter}^{(2)}=\int d^4x\sqrt{-g}\Big[-\frac{1}{2}R^2g^{\mu\nu}\partial_\mu\delta\theta\,\partial_\nu\delta\theta-\frac{1}{2}g^{\mu\nu}\partial_\mu\sigma\,\partial_\nu\sigma-R\dot{\theta}_0\,\sigma\partial^0\delta\theta-\frac{1}{2}(V''(\chi_0)-\dot{\theta}_0^2)\sigma^2\Big]\,. \tag{21}$$

The mixing coupling $\sigma\delta\dot{\theta}$ converts the σ^3 coupling, for example, into three point functions of $\delta\theta$, and hence this model can potentially give a large non-Gaussianities. Furthermore, it is known that the squeezed limit of scalar three point functions is sensitive to the mass of σ:

$$\lim_{k_3/k_1=k_3/k_2=\kappa\to0}\langle\zeta_{\mathbf{k}_1}\zeta_{\mathbf{k}_2}\zeta_{\mathbf{k}_3}\rangle\propto\kappa^{-3/2-\nu}k_1^{-6}\,, \tag{22}$$

---

[4] Strictly speaking, the name of this gauge may be inadequate because there are still tensor fluctuations and hence the spatial hypersurface is not exactly flat.

where $\nu = \sqrt{\frac{9}{4} - \frac{m_\sigma^2}{H^2}}$ and $m_\sigma^2 = V''(\chi_0) - R\dot{\theta}_0^2 < \frac{9}{4}H$. As this simple model implies, massive scalar fields of Hubble scale mass can cause a non-trivial behavior of non-Gaussianities. In the following sections we discuss more general setting for quasi-single field inflation using the effective field theory approach.

III. MOST GENERIC ACTION OF QUASI-SINGLE FIELD INFLATION

In this section, we construct the most generic action of quasi-single field inflation using the effective field theory approach. After constructing the action in the unitary gauge first, we derive the action for the Goldstone boson π and discuss its decoupling regime. Relations between our approach and models in the literatures are also discussed.

A. Action in the unitary gauge

In the unitary gauge, the relevant degrees of freedom in quasi-single field inflation are three physical modes of graviton and an additional scalar field σ. The typical mass of σ is supposed to be of the order of the Hubble scale during the inflationary era. In this subsection, we construct the most generic action invariant under the time-dependent spatial diffeomorphism from graviton and the scalar field σ up to the third order fluctuations. Here it should be noticed that the action constructed in this section can be applied to any two field models because no conditions on σ are imposed.[5]

Extending the procedures in [27] to our case, the most general action invariant under the time-dependent spatial diffeomorphism is given by

$$S = \int d^4x \sqrt{-g}\, F(R_{\mu\nu\rho\sigma}, g^{00}, K_{\mu\nu}, \nabla_\mu, t, \sigma)\,, \tag{23}$$

and it is expanded around the given FRW background as

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} M_{\rm Pl}^2 R + M_{\rm Pl}^2 \dot{H} g^{00} - M_{\rm Pl}^2 (3H^2 + \dot{H}) + F^{(2)}(\delta g^{00}, \sigma, \delta K_{\mu\nu}, \delta R_{\mu\nu\rho\sigma}; \delta_\mu^0, g_{\mu\nu}, g^{\mu\nu}, \nabla_\mu, t) \right], \tag{24}$$

where all the free indices inside the functions F and $F^{(2)}$ must be again upper 0's and $F^{(2)}$ starts with quadratic terms of the arguments δg^{00}, σ, $\delta K_{\mu\nu}$, and $\delta R_{\mu\nu\rho\sigma}$.[6] Then, let us write down possible terms in the action up to the third order fluctuations. Schematically, we write the action in the following way:

$$S = S_{\rm grav} + S_\sigma + S_{\rm mix}\,, \tag{26}$$

where the first term $S_{\rm grav}$ in the right hand side denotes terms constructed from δg^{00}, $\delta K_{\mu\nu}$, and $\delta R_{\mu\nu\rho\sigma}$, the second term S_σ denotes those only from σ, and the last term $S_{\rm mix}$ denotes those mixing the graviton fluctuations and σ. As discussed in [27], the first term $S_{\rm grav}$ can be expanded as

$$S_{\rm grav} = \int d^4x \sqrt{-g} \left[\frac{1}{2} M_{\rm Pl}^2 R + M_{\rm Pl}^2 \dot{H}(t) g^{00} - M_{\rm Pl}^2 \left(3H^2(t) + \dot{H}(t)\right) + \frac{M_2^4(t)}{2!}(\delta g^{00})^2 + \frac{M_3^4(t)}{3!}(\delta g^{00})^3 + \ldots \right], \tag{27}$$

---

[5] For example, we do not require the shift symmetry of σ, $\sigma \to \sigma + \text{constant}$, which is assumed in multi-field inflation [28]. The mass of σ is not necessarily of order Hubble scale so that the action constructed in this section can be applied not only for quasi-single field inflation but also inflation models with an additional heavy scalar.

[6] In general, sums of terms linear in the fluctuations can be practically second order. For example, let us consider the term $\int d^4x \sqrt{-g}\left[f_1(t)\sigma + f_2(t)\partial^0\sigma\right]$. Although this kind of action seems to be first order apparently, it turns out to be second order after taking into account the equation of motion for σ: $f_1(t) + \dot{f}_2(t) + 3Hf_2(t) = 0$. Then, the function $F^{(2)}$ seems to contain such a combination of linear order terms. However, using the relation

$$\begin{aligned}\int d^4x \sqrt{-g} f(t) \nabla^0(\ldots) &= -\int d^4x \sqrt{-g} f(t) \sqrt{-g^{00}} n^\mu \partial_\mu(\ldots) = \int d^4x \sqrt{-g} \Big(- g^{00} \dot{f}(t) - \frac{1}{2} f(t) \partial^0 \ln(-g^{00}) + f(t)\sqrt{-g^{00}} K^\mu_\mu \Big)(\ldots) \\ &= \int d^4x \sqrt{-g} \Big(\dot{f}(t) + 3Hf(t) + \mathcal{O}(\delta g^{00}, \delta K^\mu_\mu) \Big)(\ldots)\,, \end{aligned} \tag{25}$$

we can rewrite it into the second and higher order terms in σ, δg^{00}, and δK^μ_μ. Similar discussions hold for more general cases and we conclude that $F^{(2)}$ starts with quadratic terms of δg^{00}, σ, $\delta K_{\mu\nu}$, and $\delta R_{\mu\nu\rho\sigma}$. See also appendix B in [27].

where the dots stand for terms of higher order in the fluctuations or with more derivatives. When we rewrite the action in the unitary gauge in terms of the Goldstone boson π, the terms displayed in (27) are described by π and its first order derivatives. In this paper, we consider the action up to the same order in derivatives of π and σ.

Let us first construct the second order action. The second order action $S_\sigma^{(2)}$ containing σ and its first order derivative can be written generally as

$$S_\sigma^{(2)} = \int d^4x\sqrt{-g}\Big[-\frac{\alpha_1(t)}{2}g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma + \frac{\alpha_2(t)}{2}(\partial^0\sigma)^2 - \frac{\alpha_3(t)}{2}\sigma^2 + \alpha_4(t)\sigma\partial^0\sigma\Big]\,, \tag{28}$$

where we note that terms such as $\sigma(\partial^0)^2\sigma$ can be absorbed into other terms up to higher order fluctuations by integrating by parts. Here the second term leads to a non-trivial sound speed c_σ of σ given by $c_\sigma^2 = \alpha_1/(\alpha_1+\alpha_2)$. The second order mixing $S_{\rm mix}^{(2)}$ is generally given by

$$S_{\rm mix}^{(2)} = \int d^4x\sqrt{-g}\Big[\overline{\beta}_1(t)\delta g^{00}\sigma + \overline{\beta}_2(t)\delta g^{00}\partial^0\sigma + \beta_3(t)\delta K^\mu_\mu\sigma\Big]\,. \tag{29}$$

It is convenient to note the relation

$$\int d^4x\sqrt{-g}f(t)\delta K^\mu_\mu(\dots) = \int d^4x\sqrt{-g}\Big[f(t)\partial^0(\dots) - \Big(\dot{f}(t)+3Hf(t)\Big)(\dots) + \frac{\dot{f}(t)}{2}\delta g^{00}(\dots) + \frac{f(t)}{2}\delta g^{00}\partial^0(\dots) + \dots\Big]\,, \tag{30}$$

which can be obtained using (25) in footnote 6 twice. Here the last dots stand for higher order terms in the fluctuations, which can be written using δg^{00}, $\partial^0\delta g^{00}$, and δK^μ_μ. Using this relation (30), $S_{\rm mix}^{(2)}$ in (29) is rewritten as

$$S_{\rm mix}^{(2)} = \int d^4x\sqrt{-g}\Big[\beta_1(t)\delta g^{00}\sigma + \beta_2(t)\delta g^{00}\partial^0\sigma + \beta_3(t)\partial^0\sigma - (\dot{\beta}_3(t)+3H\beta_3(t))\sigma\Big]\,, \tag{31}$$

where $\beta_1 = \overline{\beta}_1 + \dot{\beta}_3/2$, $\beta_2 = \overline{\beta}_2 + \beta_3/2$, and we dropped higher order terms constructed from σ, δg^{00}, $\partial^0\delta g^{00}$, and δK^μ_μ. In the following, we employ Eq. (31) as a definition of the second order mixing action $S_{\rm mix}^{(2)}$.

The third order action $S_\sigma^{(3)}$ of σ is generally given by

$$S_\sigma^{(3)} = \int d^4x\sqrt{-g}\Big[\gamma_1(t)\sigma^3 + \gamma_2(t)\sigma^2\partial^0\sigma + \gamma_3(t)\sigma(\partial^0\sigma)^2 + \gamma_4(t)(\partial^0\sigma)^3 + \gamma_5(t)\,\sigma\partial_\mu\sigma\partial^\mu\sigma + \gamma_6(t)\partial^0\sigma\partial_\mu\sigma\partial^\mu\sigma\Big]\,, \tag{32}$$

and the third order mixing $S_{\rm mix}^{(3)}$ is given by

$$\begin{aligned}S_{\rm mix}^{(3)} = \int d^4x\sqrt{-g}\Big[&\overline{\gamma}_1(t)\delta g^{00}\,\sigma^2 + \overline{\gamma}_2(t)\delta g^{00}\,\sigma\partial^0\sigma + \overline{\gamma}_3(t)\delta g^{00}(\partial^0\sigma)^2 + \overline{\gamma}_4(t)\partial^0\delta g^{00}\sigma\partial^0\sigma\\ &+ \overline{\gamma}_5(t)\,\delta g^{00}\partial_\mu\sigma\partial^\mu\sigma + \overline{\gamma}_6(t)(\delta g^{00})^2\sigma + \overline{\gamma}_7(t)(\delta g^{00})^2\partial^0\sigma\Big]\,.\end{aligned} \tag{33}$$

Here, it may be wondered if the terms such as $\delta K^\mu_\mu\,\sigma^2$, $\delta K^\mu_\mu\,\sigma\partial^0\sigma$, $\delta R^{00}\sigma^2$, and $\delta K^\mu_\mu\delta g^{00}\,\sigma$ can appear at the same order in derivatives. However, they can be absorbed into other third order terms in (33) and the second order terms in (28) and (31) by integrating by parts as we did to rewrite (29) into (31). The term proportional to $\delta R\sigma^2$ can also appear but such a term vanishes in the decoupling limit, so we do not consider it here for simplicity.

To summarize, the most generic action in the unitary gauge can be written up to the third order fluctuations as follows:

$$S = S_{\rm grav} + S_\sigma^{(2)} + S_{\rm mix}^{(2)} + S_\sigma^{(3)} + S_{\rm mix}^{(3)}\,, \tag{34}$$

where $S_{\rm grav}$, $S_\sigma^{(2)}$, $S_{\rm mix}^{(2)}$, $S_\sigma^{(3)}$, and $S_{\rm mix}^{(3)}$ are defined in (27), (28), (31), (32), and (33).

In the last of this subsection, we discuss the normalization of σ. Using the field redefinition of σ, we can drop some terms in the action and consider the following three normalizations.

a. Normalization 1 : $\alpha_1+\alpha_2=1$. Let us first consider the kinetic term of σ. The second order action $S_\sigma^{(2)}$ of σ can be expanded up to the second order fluctuations as

$$\int d^4x\,a^3\left[\frac{\alpha_\sigma^2}{2}\Big(\dot\sigma^2-c_\sigma^2\frac{(\partial_i\sigma)^2}{a^2}-\frac{\alpha_3-3H\alpha_4-\dot\alpha_4}{\alpha_1+\alpha_2}\sigma^2\Big)\right], \tag{35}$$

where the normalization factor α_σ is defined as $\alpha_\sigma^2=\alpha_1+\alpha_2$. Although the factor α_σ is time-dependent in general, it can be taken unity by redefining σ as $\tilde\sigma=\alpha_\sigma\sigma$. Since the derivative of σ can be written as

$$\partial_\mu\sigma=\alpha_\sigma^{-1}\partial_\mu\tilde\sigma-\delta_\mu^0\,\frac{\dot\alpha_\sigma}{\alpha_\sigma^2}\tilde\sigma\,, \tag{36}$$

the action still takes the form (28) after the redefinition.

b. Normalization 2 : $\alpha_4=\beta_3=\gamma_2=0$. The action S in Eq. (34) contains the following terms:

$$\int d^4x\sqrt{-g}\Big[-\frac{1}{2}f_1^2(t,\sigma)g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma+f_1(t,\sigma)f_2(t,\sigma)\partial^0\sigma\Big], \tag{37}$$

where $f_1(t,\sigma)$ and $f_2(t,\sigma)$ are functions of t and σ but do not contain derivatives of σ. They are expanded in σ as $f_1^2=\alpha_1-2\gamma_5\sigma+\mathcal{O}(\sigma^2)$ and $f_1f_2=\beta_3+\alpha_4\sigma+\gamma_2\sigma^2+\mathcal{O}(\sigma^3)$. The function f_2 can be set to zero by redefining σ such that

$$f_1(t,\sigma)\partial_\mu\sigma-\delta_\mu^0\,f_2(t,\sigma)=h(t,\sigma)\partial_\mu\tilde\sigma\,, \tag{38}$$

where $h(t,\sigma)$ is some function of t and σ but do not depend on derivatives of σ. After the redefinition, the action still takes the form (34) and the second order action containing σ simplifies as

$$S_\sigma^{(2)}+S_{\rm mix}^{(2)}=\int d^4x\sqrt{-g}\Big[-\frac{\alpha_1(t)}{2}g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma+\frac{\alpha_2(t)}{2}(\partial^0\sigma)^2-\frac{\alpha_3(t)}{2}\sigma^2+\beta_1(t)\delta g^{00}\sigma+\beta_2(t)\delta g^{00}\partial^0\sigma\Big]. \tag{39}$$

c. Normalization 3 : $\alpha_1=1,\ \gamma_5=0$. The action S in Eq. (34) contains the following term:

$$\int d^4x\sqrt{-g}\Big[-\frac{1}{2}f^2(t,\sigma)g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma\Big], \tag{40}$$

where $f(t,\sigma)$ is a function of t and σ, and does not contain derivatives of σ. The function is expanded in σ as $f^2(t,\sigma)=\alpha_1(t)-2\gamma_5(t)\sigma+\mathcal{O}(\sigma^2)$. When α_2 is small enough, in other words, the sound speed of σ is almost unity, it would be convenient to take $f(t,\sigma)$ to be unity. Let us define $\tilde\sigma$ as $\tilde\sigma=F(\sigma,t)$ such that $\partial F/\partial\sigma=f$. Since the derivative of $\tilde\sigma$ is given by

$$\partial_\mu\tilde\sigma=f\partial_\mu\sigma+\delta_\mu^0\frac{\partial F}{\partial t}\,, \tag{41}$$

the action can be rewritten as

$$\int d^4x\sqrt{-g}\Big[-\frac{1}{2}g^{\mu\nu}\partial_\mu\tilde\sigma\partial_\nu\tilde\sigma+\frac{\partial F}{\partial t}\partial^0\tilde\sigma-\frac{1}{2}g^{00}\left(\frac{\partial F}{\partial t}\right)^2\Big], \tag{42}$$

which still takes the form (34). Here F should be regarded as a function of $\tilde\sigma$ and t. It also should be noticed that, though we can set the function f to be unity, the terms proportional to $\partial^0\sigma$ and g^{00} appear.

B. Action for the Goldstone boson and the decoupling limit

In this subsection we construct the action for the Goldstone boson π and discuss its decoupling regime. As in the last section, we perform the time diffeomorphism (10). Practically, it is realized by the following replacements:

$$\begin{aligned}&g^{00}\to g^{00}+2\partial^0\pi+\partial_\mu\pi\partial^\mu\pi\,,\quad \partial^0\sigma\to\partial^0\sigma+\partial_\mu\pi\,\partial^\mu\sigma\,,\\&\sigma\to\sigma\,,\quad f(t)\to f(t+\pi)\,,\quad \int d^4x\sqrt{-g}\to\int d^4x\sqrt{-g}\,.\end{aligned} \tag{43}$$

With these replacements, $S_{\rm grav}$, $S_\sigma^{(2)}$, and $S_{\rm mix}^{(2)}$ are rewritten as

$$
\begin{aligned}
S_{\rm grav} = \int d^4x\sqrt{-g}\bigg[&\frac{1}{2}M_{\rm Pl}^2 R + M_{\rm Pl}^2\dot{H}(t+\pi)\left(g^{00}+2\partial^0\pi+\partial_\mu\pi\partial^\mu\pi\right) - M_{\rm Pl}^2\Big(3H^2(t+\pi)+\dot{H}(t+\pi)\Big) \\
&+\frac{M_2^4(t+\pi)}{2!}\left(\delta g^{00}+2\partial^0\pi+\partial_\mu\pi\partial^\mu\pi\right)^2+\frac{M_3^4(t+\pi)}{3!}\left(\delta g^{00}+2\partial^0\pi+\partial_\mu\pi\partial^\mu\pi\right)^3\bigg]\,, \qquad (44)
\end{aligned}
$$

$$
S_\sigma^{(2)} = \int d^4x\sqrt{-g}\Big[-\frac{\alpha_1(t+\pi)}{2}\partial_\mu\sigma\partial^\mu\sigma+\frac{\alpha_2(t+\pi)}{2}\left(\partial^0\sigma+\partial_\mu\pi\partial^\mu\sigma\right)^2-\frac{\alpha_3(t+\pi)}{2}\sigma^2+\alpha_4(t+\pi)\sigma(\partial^0\sigma+\partial_\mu\pi\,\partial^\mu\sigma)\Big]\,, \qquad (45)
$$

$$
\begin{aligned}
S_{\rm mix}^{(2)} = \int d^4x\sqrt{-g}\Big[&\beta_1(t+\pi)\left(\delta g^{00}+2\partial^0\pi+\partial_\mu\pi\partial^\mu\pi\right)\sigma+\beta_2(t+\pi)\left(\delta g^{00}+2\partial^0\pi+\partial_\mu\pi\partial^\mu\pi\right)\left(\partial^0\sigma+\partial_\mu\pi\partial^\mu\sigma\right) \\
&+\beta_3(t+\pi)(\partial^0\sigma+\partial_\mu\pi\partial^\mu\sigma)-\Big(\dot\beta_3(t+\pi)+3H(t+\pi)\beta_3(t+\pi)\Big)\sigma\Big]\,. \qquad (46)
\end{aligned}
$$

The third order actions $S_\sigma^{(3)}$ and $S_{\rm mix}^{(3)}$ can be obtained in a similar way.

In order to discuss the decoupling regime of the action, we first clarify in which regime graviton fluctuations become irrelevant to tree-level three point functions of π. For this purpose, let us take the spatially flat gauge (18) and use the ADM decomposition:

$$
ds^2=-(N^2-N_iN^i)dt^2+2N_idx^idt+a^2(e^\gamma)_{ij}\,dx^idx^j \quad \text{with} \quad \gamma_{ii}=\partial_i\gamma_{ij}=0\,. \qquad (47)
$$

Here and in what follows we use the spatial metric $h_{ij}=a^2(e^\gamma)_{ij}$ and its inverse $h^{ij}=a^{-2}(e^{-\gamma})_{ij}$ to raise or lower the indices of N^i. The inverse metric $g^{\mu\nu}$ are written in terms of N, N^i, and h_{ij} as

$$
g^{00}=-\frac{1}{N^2}\,,\quad g^{0i}=g^{i0}=\frac{N^i}{N^2}\,,\quad g^{ij}=h^{ij}-\frac{N^iN^j}{N^2}\,. \qquad (48)
$$

In this gauge, there are no second order mixing terms of π and γ_{ij} because γ_{ij} has two spatial indices and is transverse-traceless. Then, the tensor fluctuation γ_{ij} does not contribute to tree-level three point functions of π. Therefore, possible contributions of graviton fluctuations come only from the auxiliary fields $\delta N=N-1$ and N^i. As discussed in [30], it is sufficient for the calculation of three-point functions to solve the constraints up to first order. Expanding the actions (44)-(46) up to the second order in π, δN, and N^i,

$$
\begin{aligned}
S_{\rm grav} = \int d^4x\,a^3\bigg[&-M_{\rm Pl}^2(3H^2+c_\pi^{-2}\dot H)\delta N^2-2M_{\rm Pl}^2H\delta N\partial_iN^i+M_{\rm Pl}^2\frac{1}{4}N^i\partial_i\partial_jN^j-M_{\rm Pl}^2\frac{1}{4}N^i\partial^2N^i \\
&-\frac{M_{\rm Pl}^2\dot H}{c_\pi^2}\Big(\dot\pi^2-c_\pi^2\frac{(\partial_i\pi)^2}{a^2}\Big)-3M_{\rm Pl}^2\dot H^2\pi^2+M_{\rm Pl}^2(2c_\pi^{-2}\dot H\dot\pi-6H\dot H\pi)\delta N+2M_{\rm Pl}^2\dot HN^i\partial_i\pi\bigg]\,, \qquad (49)
\end{aligned}
$$

$$
S_\sigma^{(2)} = \int d^4x\,a^3\Big[\alpha_\sigma^2\Big(\dot\sigma^2-c_\sigma^2\frac{(\partial_i\sigma)^2}{a^2}\Big)-\frac{\alpha_3}{2}\sigma^2-\alpha_4\sigma\dot\sigma\Big]\,, \qquad (50)
$$

$$
\begin{aligned}
S_{\rm mix}^{(2)} = \int d^4x\,a^3\Big[&2\beta_1\left(\delta N-\dot\pi\right)\sigma-2\beta_2\left(\delta N-\dot\pi\right)\dot\sigma-\delta N\Big(-\beta_3\dot\sigma+\dot\beta_3\sigma+3H\beta_3\sigma\Big)+\beta_3N^i\partial_i\sigma \\
&+\beta_3\Big(-\dot\pi\dot\sigma+\frac{\partial_i\pi\partial_i\sigma}{a^2}\Big)+\dot\beta_3\dot\pi\sigma-3\dot H\beta_3\pi\sigma\Big]\,, \qquad (51)
\end{aligned}
$$

the constraints are solved up to first order as follows:

$$
\begin{aligned}
&\delta N=-\frac{\dot H}{H}\pi-\frac{\beta_3}{2M_{\rm Pl}^2H}\sigma\,,\quad N^i=a^{-2}\partial_i\psi \\
&\qquad \text{with} \quad \psi=a^2\partial^{-2}\Big(c_\pi^{-2}\frac{\dot H}{H^2}(\dot H\pi+H\dot\pi)+\frac{\beta_1}{M_{\rm Pl}^2H}\sigma-\frac{\beta_2}{M_{\rm Pl}^2H}\dot\sigma+\frac{\beta_3}{2M_{\rm Pl}^2H}\Big(c_\pi^{-2}\frac{\dot H}{H}\sigma+\dot\sigma\Big)-\frac{\dot\beta_3}{2M_{\rm Pl}^2H}\sigma\Big)\,. \qquad (52)
\end{aligned}
$$

Here the sound speed c_π^2 of π are defined as $c_\pi^2=\dot HM_{\rm Pl}^2/(\dot HM_{\rm Pl}^2-2M_2^4)$. The factors $c_\sigma^2=\alpha_1/(\alpha_1+\alpha_2)$ and $\alpha_\sigma^2=\alpha_1+\alpha_2$ are the sound speed and the normalization factor of σ, respectively. Using the canonical normalization

$$
\pi_c\sim M_{\rm Pl}(-\dot H)^{1/2}c_\pi^{-1}\pi\,,\quad \sigma_c\sim\alpha_\sigma\sigma\,,\quad \delta N_c\sim M_{\rm Pl}\delta N\,,\quad N_c^i\sim M_{\rm Pl}N^i\,, \qquad (53)
$$

and redefining the coupling constants β_1, β_2, and β_3 correspondingly as

$$\beta_1^c \sim \frac{c_\pi}{\alpha_\sigma M_{\rm Pl}(-\dot{H})^{1/2}}\beta_1\,, \quad \beta_2^c \sim \frac{c_\pi}{\alpha_\sigma M_{\rm Pl}(-\dot{H})^{1/2}}\beta_2\,, \quad \beta_3^c \sim \frac{c_\pi}{\alpha_\sigma M_{\rm Pl}(-\dot{H})^{1/2}}\beta_3\,, \tag{54}$$

we rewrite the constraints (52) as

$$\delta N_c \sim \tilde{\epsilon}^{1/2}\Big(c_\pi^2\pi_c - \frac{1}{2}\beta_3^c\,\sigma_c\Big)\,, \tag{55}$$

$$N_c^i \sim \tilde{\epsilon}^{1/2}\frac{\partial_i}{\partial^2}\left(-\dot{\pi}_c + \frac{1}{2}\tilde{\eta}H\pi_c + \beta_1^c\sigma_c - \beta_2^c\Big(\dot{\sigma}_c - \frac{\dot{\alpha}_\sigma}{\alpha_\sigma}\sigma\Big) - \frac{1}{2}\beta_3^c\dot{\sigma}_c + \frac{1}{2}\beta_3^c\sigma_c\Big(-2\frac{\dot{\alpha}_\sigma}{\alpha_\sigma}\sigma + (c_\pi^2-1)\tilde{\epsilon}H - \frac{\tilde{\eta}}{2}H\Big) - \frac{1}{2}\dot{\beta}_3^c\sigma_c\right)\,, \tag{56}$$

where we have defined $\tilde{\epsilon} = -c_\pi^{-2}\dfrac{\dot{H}}{H^2}$ and $\tilde{\eta} = \dfrac{\dot{\tilde{\epsilon}}}{\tilde{\epsilon}H}$ in analogy with usual slow-roll parameters ϵ and η. It is manifest that δN_c and N_c^i are suppressed by the parameter $\tilde{\epsilon}^{1/2}$ and contributions from δN_c and N_c^i become irrelevant in the limit $\tilde{\epsilon} \to 0$.[7] In this limit, tree-level three point functions of π are determined by the following action in the decoupling limit:

$$\begin{aligned} S_{\rm grav} = \int d^4x\,a^3\Big[&-\frac{M_{\rm Pl}^2\dot{H}}{c_\pi^2}\Big(\dot{\pi}^2 - c_\pi^2\frac{(\partial_i\pi)^2}{a^2}\Big) - M_{\rm Pl}^2\dot{H}(c_\pi^{-2}-1)\Big(\dot{\pi}^3 - \dot{\pi}\frac{(\partial_i\pi)^2}{a^2}\Big) - \frac{4M_3^4}{3}\dot{\pi}^3 \\ &- 3M_{\rm Pl}^2\dot{H}^2\pi^2 - \partial_t\Big(\frac{M_{\rm Pl}^2\dot{H}}{c_\pi^2}\Big)\dot{\pi}^2\pi + M_{\rm Pl}^2\ddot{H}\frac{(\partial_i\pi)^2}{a^2}\pi - 3M_{\rm pl}^2\dot{H}\ddot{H}\pi^3\Big]\,, \end{aligned} \tag{57}$$

$$\begin{aligned} S_\sigma^{(2)} = \int d^4x\,a^3\Big[&\frac{\alpha_\sigma^2}{2}\Big(\dot{\sigma}^2 - c_\sigma^2\frac{(\partial_i\sigma)^2}{a^2} - \frac{\alpha_3 - 3H\alpha_4 - \dot{\alpha}_4}{\alpha_\sigma^2}\sigma^2\Big) + \alpha_\sigma^2(1-c_\sigma^2)\Big(\dot{\pi}\dot{\sigma}^2 - \dot{\sigma}\frac{(\partial_i\pi\partial_i\sigma)}{a^2}\Big) \\ &- \alpha_4\sigma\Big(\dot{\pi}\dot{\sigma} - \frac{\partial_i\pi\partial_i\sigma}{a^2}\Big) + \frac{\dot{(\alpha_\sigma^2)}}{2}\pi\dot{\sigma}^2 - \frac{\dot{(\alpha_\sigma^2c_\sigma^2)}}{2}\pi\frac{(\partial_i\sigma)^2}{a^2} - \frac{\dot{\alpha}_3}{2}\sigma^2\pi - \dot{\alpha}_4\pi\sigma\dot{\sigma}\Big]\,, \end{aligned} \tag{58}$$

$$\begin{aligned} S_{\rm mix}^{(2)} = \int d^4x\,a^3\Big[&(-2\beta_1 + \dot{\beta}_3)\dot{\pi}\sigma + (2\beta_2 - \beta_3)\dot{\pi}\dot{\sigma} + \beta_3\frac{\partial_i\pi\partial_i\sigma}{a^2} - 3\dot{H}\beta_3\pi\sigma - \beta_1\Big(\dot{\pi}^2 - \frac{(\partial_i\pi)^2}{a^2}\Big)\sigma + (-2\dot{\beta}_1 + \ddot{\beta}_3)\pi\dot{\pi}\sigma \\ &+ 3\beta_2\dot{\pi}^2\dot{\sigma} - 2\beta_2\dot{\pi}\frac{\partial_i\pi\partial_i\sigma}{a^2} - \beta_2\frac{(\partial_i\pi)^2}{a^2}\dot{\sigma} + (2\dot{\beta}_2 - \dot{\beta}_3)\pi\dot{\pi}\dot{\sigma} + \dot{\beta}_3\pi\Big(\frac{\partial_i\pi\partial_i\sigma}{a^2}\Big) - 3\left(\dot{H}\dot{\beta}_3 + \frac{1}{2}\ddot{H}\beta_3\right)\pi^2\sigma\Big]\,, \end{aligned} \tag{59}$$

$$S_\sigma^{(3)} = \int d^4x\,a^3\Big[\Big(\gamma_1 + H\gamma_2 + \frac{1}{3}\dot{\gamma}_2\Big)\sigma^3 + (\gamma_3 - \gamma_5)\sigma\dot{\sigma}^2 + (-\gamma_4 + \gamma_6)\dot{\sigma}^3 + \gamma_5\,\sigma\frac{(\partial_i\sigma)^2}{a^2} - \gamma_6\dot{\sigma}\frac{(\partial_i\sigma)^2}{a^2}\Big]\,, \tag{60}$$

$$S_{\rm mix}^{(3)} = \int d^4x\,a^3\Big[-2\overline{\gamma}_1\dot{\pi}\sigma^2 + 2\overline{\gamma}_2\dot{\pi}\sigma\dot{\sigma} + 2(-\overline{\gamma}_3 + \overline{\gamma}_5)\dot{\pi}\dot{\sigma}^2 - 2\overline{\gamma}_4\ddot{\pi}\sigma\dot{\sigma} - 2\overline{\gamma}_5\dot{\pi}\frac{(\partial_i\sigma)^2}{a^2} + 4\overline{\gamma}_6\dot{\pi}^2\sigma - 4\overline{\gamma}_7\dot{\pi}^2\dot{\sigma}\Big]\,. \tag{61}$$

It should be noticed that non-trivial cubic interactions appear generically when the sound speed c_σ of σ is small, α_4 is non-zero, or mixing couplings β_1 and β_2 exist as well as the sound speed c_π of π is small.

C. Examples

In the last of this section, we clarify the relation between our approach and models in the literatures. For this purpose, we first discuss the original model of quasi-single field inflation [21], and then, we investigate the effects of heavy particles during inflation. In the last of this subsection, a class of two field models will be considered.

---

[7] To be precise, we need to assume that $\tilde{\epsilon}$ is small enough to vanish when multiplied by other parameters in the calculation. For example, we need to assume that $\tilde{\epsilon}\frac{H}{E} \ll 1$.

1. *Original model discussed by Chen and Wang*

As was reviewed in section II, the original model [21] of quasi-single field inflation is described by the following matter action:

$$S_{\rm matter}=\int d^4x\sqrt{-g}\Big[-\frac{1}{2}(\tilde{R}+\chi)^2g^{\mu\nu}\partial_\mu\theta\partial_\nu\theta-\frac{1}{2}g^{\mu\nu}\partial_\mu\chi\partial_\nu\chi-V_{\rm sr}(\theta)-V(\chi)\Big]\,. \tag{62}$$

The homogeneous backgrounds are given by $\theta=\theta_0(t)$ and $\chi=\chi_0$ (constant), which leads to the action in the unitary gauge $\delta\theta=\theta-\theta_0=0$,

$$\begin{aligned}S_{\rm matter}&=\int d^4x\sqrt{-g}\Big[-\frac{1}{2}(R+\sigma)^2g^{00}\dot{\theta}_0^2-\frac{1}{2}g^{\mu\nu}\partial_\mu\sigma\,\partial_\nu\sigma-V_{\rm sr}(\theta_0)-V(\chi_0+\sigma)\Big]\\&=\int d^4x\sqrt{-g}\Big[-\frac{1}{2}R^2\dot{\theta}_0^2\,g^{00}-(V_{\rm sr}(\theta_0)+V(\chi_0))-\frac{1}{2}g^{\mu\nu}\partial_\mu\sigma\,\partial_\nu\sigma-\frac{1}{2}\Big(V''(\chi_0)-\dot{\theta}_0^2\Big)\sigma^2\\&\qquad\qquad-R\,\dot{\theta}_0^2\,\delta g^{00}\sigma-\frac{V'''(\chi_0)}{3!}\sigma^3-\frac{1}{2}\dot{\theta}_0^2\,\delta g^{00}\sigma^2+\mathcal{O}(\sigma^4)\Big]\,,\end{aligned} \tag{63}$$

where $R=\tilde{R}+\chi_0$, $\sigma=\chi-\chi_0$ and we used the background equations of motion. Using the equations of motion (20), the action (63) can be written in terms of the Hubble parameter H as

$$\begin{aligned}S_{\rm matter}=\int d^4x\sqrt{-g}\Big[&M_{\rm Pl}^2\dot{H}g^{00}-M_{\rm Pl}^2(3H^2+\dot{H})-\frac{1}{2}g^{\mu\nu}\partial_\mu\sigma\,\partial_\nu\sigma-\frac{1}{2}\Big(V''(\chi_0)+\frac{2M_{\rm Pl}^2\dot{H}}{R^2}\Big)\sigma^2\\&+\frac{2M_{\rm Pl}^2\dot{H}}{R}\,\delta g^{00}\sigma-\frac{V'''(\chi_0)}{3!}\sigma^3+\frac{M_{\rm Pl}^2\dot{H}}{R^2}\,\delta g^{00}\sigma^2+\mathcal{O}(\delta\sigma^4)\Big]\,,\end{aligned} \tag{64}$$

which corresponds to the following parameters in our framework,

$$\alpha_1=1\,,\quad\alpha_3=V''(\chi_0)+\frac{2M_{\rm Pl}^2\dot{H}}{R^2}\,,\quad\beta_1=\frac{2M_{\rm Pl}^2\dot{H}}{R}\,,\quad\gamma_1=-\frac{1}{3!}V'''(\chi_0)\,,\quad\overline{\gamma}_1=\frac{M_{\rm Pl}^2\dot{H}}{R^2}\,,\quad(\,\text{others}\,)=0\,. \tag{65}$$

2. *Effects of heavy particles*

Recently, it is argued that the existence of heavy particles can cause a non-trivial sound speed of effective single field inflation [13–20]. As was mentioned earlier, our framework is also applicable for such inflation models with heavy particles. In the following, we briefly show that it gives a simple explanation for the effective sound speed.

As a simplest example, let us consider the following quadratic action:

$$S=\int d^4x\sqrt{-g}\left[\frac{1}{2}M_{\rm Pl}^2R+M_{\rm Pl}^2\dot{H}g^{00}-M_{\rm Pl}^2(3H^2+\dot{H})-\frac{1}{2}g^{\mu\nu}\partial_\mu\sigma\,\partial_\nu\sigma-\frac{m^2}{2}\sigma^2+\beta\,\delta g^{00}\sigma\right]\,. \tag{66}$$

Here m is the mass of σ and β is the mixing coupling between the adiabatic mode and the massive particle σ. We assume that time-dependence of m and β is negligible, and the mass of σ is much larger than the Hubble scale during inflation, $m\gg H$. In such a regime, the kinetic term of σ becomes irrelevant and the dynamics is determined by

$$S\sim\int d^4x\sqrt{-g}\left[\frac{1}{2}M_{\rm Pl}^2R+M_{\rm Pl}^2\dot{H}g^{00}-M_{\rm Pl}^2(3H^2+\dot{H})-\frac{m^2}{2}\Big(\sigma-\frac{\beta}{m^2}\delta g^{00}\Big)^2+\frac{\beta^2}{2m^2}(\delta g^{00})^2\right]\,, \tag{67}$$

which implies that the perturbation σ quickly responds to the variation of the adiabatic mode δg^{00}. Integrating out the massive particle σ, we obtain the following effective action for single-field inflation:

$$S=\int d^4x\sqrt{-g}\left[\frac{1}{2}M_{\rm Pl}^2R+M_{\rm Pl}^2\dot{H}g^{00}-M_{\rm Pl}^2(3H^2+\dot{H})+\frac{\beta^2}{2m^2}(\delta g^{00})^2\right]\,. \tag{68}$$

As we discussed earlier, the last term gives the following non-trivial sound speed:

$$c_\pi^2=\frac{-\dot{H}M_{\rm Pl}^2}{-\dot{H}M_{\rm Pl}^2+2\beta^2/m^2}\,. \tag{69}$$

3. A class of two-field models

Let us then consider a class of two-field models described by the following matter action:

$$S_{\rm matter}=\int d^4x\sqrt{-g}\Big[-\frac{1}{2}\gamma_{ab}(\phi^a)g^{\mu\nu}\partial_\mu\phi^a\partial_\nu\phi^b-V(\phi^a)\Big]\quad(a=1,2)\,, \tag{70}$$

where $\gamma_{ab}(\phi^a)$ is the metric on the field space. This class of models were carefully studied in [31] and recently discussed in [15] to investigate effects of massive particles during inflation. Suppose that the trajectory of the homogeneous background fields $\phi_0^a(t)$ is on a curve $\phi^a=\bar{\phi}^a(\lambda)$, and the background fields are given by $\phi_0^a(t)=\bar{\phi}(\lambda_0(t))$. We also assume that $\dot{\lambda}>0$. Defining the coordinates (λ,σ) of the field space such that the curve $\sigma=0$ coincides with the trajectory curve, the fields $\lambda(x)$ and $\sigma(x)$ describe the adiabatic mode and the isocurvature mode, respectively. There are still many choices of the coordinates or degrees of freedom of the field redefinition. In the following, we consider two types of basis of the fields and discuss their properties in our framework.

a. Orthogonal basis We first consider the orthogonal-basis. We can always take the coordinate (λ,σ) such that

$$\gamma_{\lambda\sigma}=\gamma_{\sigma\lambda}=0\,. \tag{71}$$

By the field redefinition of λ, we further require $\gamma_{\lambda\lambda}(\lambda,\sigma=0)=1$. In this basis, the matter action (70) is given by

$$S_{\rm matter}=\int d^4x\sqrt{-g}\Big[-\frac{1}{2}\gamma_{\lambda\lambda}(\lambda,\sigma)g^{\mu\nu}\partial_\mu\lambda\partial_\nu\lambda-\frac{1}{2}\gamma_{\sigma\sigma}(\lambda,\sigma)g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma-V(\lambda,\sigma)\Big]\,, \tag{72}$$

and it can be expanded in the unitary gauge $\delta\lambda=\lambda-\lambda_0=0$ up to the third order fluctuations as

$$\begin{aligned}S_{\rm matter}=\int d^4x\sqrt{-g}\Big[&-\frac{1}{2}\dot{\lambda}_0^2\,g^{00}-\frac{1}{2}\dot{\lambda}_0^2(\gamma_{\lambda\lambda})'_0\delta g^{00}\sigma+\frac{1}{4}\dot{\lambda}_0^2(\gamma_{\lambda\lambda})''_0\sigma^2-\frac{1}{4}\dot{\lambda}_0^2(\gamma_{\lambda\lambda})''_0\delta g^{00}\sigma^2+\frac{1}{12}\dot{\lambda}_0^2(\gamma_{\lambda\lambda})'''_0\sigma^3\\&-\frac{1}{2}(\gamma_{\sigma\sigma})_0\,g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma-\frac{1}{2}(\gamma_{\sigma\sigma})'_0\,\sigma\,g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma-V_0-\frac{1}{2}V''_0\sigma^2-\frac{1}{6}V'''_0\sigma^3\Big]\,,\end{aligned} \tag{73}$$

where we have used the equation of motion for σ. In this subsection, we write derivatives of the metric and the potential evaluated at the classical value, for example, as $(\gamma_{\lambda\lambda})'_0=\partial_\sigma\gamma_{\lambda\lambda}(\lambda,\sigma)|_{\lambda=\lambda_0,\sigma=0}$ and $V'_0=\partial_\sigma V(\lambda,\sigma)|_{\lambda=\lambda_0,\sigma=0}$. Using the equations of motion for graviton,

$$M_{\rm Pl}^2\dot{H}=-\frac{1}{2}\dot{\lambda}_0^2\,,\quad M_{\rm Pl}^2(3H^2+\dot{H})=V_0\,, \tag{74}$$

we can rewrite (73) as

$$\begin{aligned}S_{\rm matter}=\int d^4x\sqrt{-g}\Big[&M_{\rm Pl}^2g^{00}\dot{H}-M_{\rm Pl}^2(3H^2+\dot{H})-\frac{1}{2}(\gamma_{\sigma\sigma})_0\,g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma-\frac{1}{2}\Big(V''_0+M_{\rm Pl}^2\dot{H}(\gamma_{\lambda\lambda})''_0\Big)\sigma^2\\&+M_{\rm Pl}^2\dot{H}(\gamma_{\lambda\lambda})'_0\delta g^{00}\sigma-\frac{1}{2}(\gamma_{\sigma\sigma})'_0\,\sigma\,g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma-\frac{1}{6}\left(V'''_0+M_{\rm Pl}^2\dot{H}(\gamma_{\lambda\lambda})'''_0\right)\sigma^3+\frac{1}{2}M_{\rm Pl}^2\dot{H}(\gamma_{\lambda\lambda})''_0\delta g^{00}\sigma^2\Big]\,,\end{aligned} \tag{75}$$

which is described in our framework as

$$\begin{aligned}&\alpha_1=(\gamma_{\sigma\sigma})_0\,,\quad\alpha_3=V''_0+M_{\rm Pl}^2\dot{H}(\gamma_{\lambda\lambda})''_0\,,\quad\beta_1=M_{\rm Pl}^2\dot{H}(\gamma_{\lambda\lambda})'_0\,,\\&\gamma_1=-\frac{1}{6}\left(V'''_0+M_{\rm Pl}^2\dot{H}(\gamma_{\lambda\lambda})'''_0\right)\,,\quad\gamma_5=-\frac{1}{2}(\gamma_{\sigma\sigma})'_0\,,\quad\overline{\gamma}_1=\frac{1}{2}M_{\rm Pl}^2\dot{H}(\gamma_{\lambda\lambda})''_0\,,\quad(\text{ others })=0\,.\end{aligned} \tag{76}$$

We notice that this basis corresponds to the second normalization of σ in section III A.

b. Canonical σ basis We then choose the coordinate (λ,σ) such that

$$\gamma_{\sigma\sigma}(\lambda,\sigma)=1\,,\quad\gamma_{\lambda\lambda}(\lambda,\sigma=0)=1\,, \tag{77}$$

where the normalization of σ is always canonical and that of λ is canonical only on the trajectory curve. In this basis, the matter action (70) is given by

$$S_{\rm matter}=\int d^4x\sqrt{-g}\Big[-\frac{1}{2}\gamma_{\lambda\lambda}(\lambda,\sigma)g^{\mu\nu}\partial_\mu\lambda\partial_\nu\lambda-\gamma_{\lambda\sigma}(\lambda,\sigma)g^{\mu\nu}\partial_\mu\lambda\partial_\nu\sigma-\frac{1}{2}g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma-V(\lambda,\sigma)\Big]\,, \tag{78}$$

and it can be expanded in the unitary gauge $\delta\lambda = \lambda - \lambda_0 = 0$ up to the third order fluctuations as

$$\begin{aligned} S_{\rm matter} = \int d^4x\sqrt{-g}\Big[& -\frac{1}{2}\dot\lambda_0^2\, g^{00} + \frac{1}{2}\dot\lambda_0^2(\gamma_{\lambda\lambda})'_0\,\sigma - \frac{1}{2}\dot\lambda_0^2(\gamma_{\lambda\lambda})'_0\delta g^{00}\sigma + \frac{1}{4}\dot\lambda_0^2(\gamma_{\lambda\lambda})''_0\sigma^2 - \frac{1}{4}\dot\lambda_0^2(\gamma_{\lambda\lambda})''_0\delta g^{00}\sigma^2 + \frac{1}{12}\dot\lambda_0^2(\gamma_{\lambda\lambda})''_0\sigma^3 \\ & - \dot\lambda_0(\gamma_{\lambda\sigma})_0\partial^0\sigma - \dot\lambda_0(\gamma_{\lambda\sigma})'_0\sigma\partial^0\sigma - \frac{1}{2}\dot\lambda_0(\gamma_{\lambda\sigma})''_0\sigma^2\partial^0\sigma - \frac{1}{2}g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma \\ & - V_0 - V'_0\sigma - \frac{1}{2}V''_0\sigma^2 - \frac{1}{6}V'''_0\sigma^3\Big]\,, \end{aligned} \tag{79}$$

where the equation of motion for σ implies $V'_0 = \frac{1}{2}\dot\lambda_0^2(\gamma_{\lambda\lambda})'_0 - \Big(3H\dot\lambda_0(\gamma_{\lambda\sigma})_0 + \ddot\lambda_0(\gamma_{\lambda\sigma})_0 + \dot\lambda_0\,\partial_t(\gamma_{\lambda\sigma})_0\Big)$. In terms of the Hubble parameter H, we can rewrite it as

$$\begin{aligned} S_{\rm matter} = \int d^4x\sqrt{-g}\Big[& M_{\rm Pl}^2 g^{00}\dot H - M_{\rm Pl}^2(3H^2+\dot H) - \sqrt{2}M_{\rm Pl}(-\dot H)^{1/2}(\gamma_{\lambda\sigma})_0\partial^0\sigma - (V'_0 + M_{\rm Pl}^2\dot H(\gamma_{\lambda\lambda})'_0)\sigma \\ & - \frac{1}{2}g^{\mu\nu}\partial_\mu\sigma\partial_\nu\sigma - \frac{1}{2}\Big(V''_0 + M_{\rm Pl}^2\dot H(\gamma_{\lambda\lambda})''_0\Big)\sigma^2 - \sqrt{2}M_{\rm Pl}(-\dot H)^{1/2}(\gamma_{\lambda\sigma})'_0\sigma\partial^0\sigma + M_{\rm Pl}^2\dot H(\gamma_{\lambda\lambda})'_0\delta g^{00}\sigma \\ & - \frac{1}{6}\left(V'''_0 + M_{\rm Pl}\dot H(\gamma_{\lambda\lambda})'''_0\right)\sigma^3 - \frac{\sqrt{2}}{2}M_{\rm Pl}(-\dot H)^{1/2}(\gamma_{\lambda\sigma})''_0\sigma^2\partial^0\sigma + \frac{1}{2}M_{\rm Pl}^2\dot H(\gamma_{\lambda\lambda})''_0\delta g^{00}\sigma^2\Big]\,, \end{aligned} \tag{80}$$

which is described in our framework as

$$\begin{aligned} & \alpha_1 = 1\,, \quad \alpha_3 = V''_0 + M_{\rm Pl}^2\dot H(\gamma_{\lambda\lambda})''_0\,, \quad \alpha_4 = -\sqrt{2}M_{\rm Pl}(-\dot H)^{1/2}(\gamma_{\lambda\sigma})'_0\,, \quad \beta_1 = M_{\rm Pl}^2\dot H(\gamma_{\lambda\lambda})'_0\,, \\ & \beta_3 = -\sqrt{2}M_{\rm Pl}(-\dot H)^{1/2}(\gamma_{\lambda\sigma})_0\,, \quad \gamma_1 = -\frac{1}{6}\left(V'''_0 + M_{\rm Pl}\dot H(\gamma_{\lambda\lambda})'''_0\right)\,, \quad \gamma_2 = -\frac{\sqrt{2}}{2}M_{\rm Pl}(-\dot H)^{1/2}(\gamma_{\lambda\sigma})''_0\,, \\ & \overline{\gamma}_1 = \frac{1}{2}M_{\rm Pl}^2\dot H(\gamma_{\lambda\lambda})''_0\,, \quad (\text{ others }) = 0\,. \end{aligned} \tag{81}$$

The above action apparently includes terms linear in σ. However, it can be easily shown that such terms turn out to be more than second order after integration by parts. We notice that this basis corresponds to the third normalization of σ in section III A.

IV. POWER SPECTRUM

In this section we calculate the power spectrum for a class of quasi-single field inflation models in our framework. In the following, we take the decoupling limit $\tilde\epsilon \to 0$ and use the action for the Goldstone boson π. It is also assumed that the background trajectory satisfies $\epsilon \ll 1$. Up to the second order fluctuations, the action for π and σ takes the following form in the decoupling limit:

$$S = \int d^4x\, a^3\Big[-\frac{M_{\rm Pl}^2\dot H}{c_\pi^2}\Big(\dot\pi^2 - c_\pi^2\frac{(\partial_i\pi)^2}{a^2}\Big) + \frac{\alpha_\sigma^2}{2}\Big(\dot\sigma^2 - c_\sigma^2\frac{(\partial_i\sigma)^2}{a^2} - m_\sigma^2\sigma^2\Big) + \tilde\beta_1\dot\pi\sigma + \tilde\beta_2\dot\pi\dot\sigma + \tilde\beta_3 c_\pi^2\frac{\partial_i\pi\partial_i\sigma}{a^2}\Big]\,, \tag{82}$$

where we have dropped sub-leading terms $\sim M_{\rm Pl}^2\dot H\pi^2$ and $\sim \dot H\pi\sigma$ in the regime $\epsilon \ll 1$, and have defined

$$m_\sigma^2 = \frac{\alpha_3 - 3H\alpha_4 - \dot\alpha_4}{\alpha_\sigma^2}\,, \quad \tilde\beta_1 = -2\beta_1 + \dot\beta_3\,, \quad \tilde\beta_2 = 2\beta_2 - \beta_3\,, \quad \tilde\beta_3 = c_\pi^{-2}\beta_3\,. \tag{83}$$

In this paper we concentrate on the case where the mixing couplings can be treated as perturbations although they can give large contributions in general. As pointed out in [21], the perturbativity of mixing couplings is justified even in the case when they are large only in a sufficiently short period of time, which is realized, for example, in the case of the sharp turning background trajectory [14–17, 19, 20].

Let us then calculate the power spectrum using the in-in formalism. Before going to concrete models, we introduce a general expression for the power spectrum up to the leading order corrections of the mixing couplings. As in the usual case, the fields π and σ are expanded in the Fourier space as

$$\pi_{\mathbf{k}} = u_k\, a_{\mathbf{k}} + u_k^*\, a_{-\mathbf{k}}^\dagger\,, \quad \sigma_{\mathbf{k}} = v_k\, b_{\mathbf{k}} + v_k^*\, b_{-\mathbf{k}}^\dagger \tag{84}$$

with the standard commutation relations

$$[a_{\mathbf{k}}, a^\dagger_{\mathbf{k}'}] = (2\pi)^3\delta^{(3)}(\mathbf{k}-\mathbf{k}')\,, \quad [b_{\mathbf{k}}, b^\dagger_{\mathbf{k}'}] = (2\pi)^3\delta^{(3)}(\mathbf{k}-\mathbf{k}')\,. \tag{85}$$

Here the mode functions u_k and v_k satisfy the equations of motion in the free theory and depend on $k = |\mathbf{k}|$:

$$\ddot{u}_k + H(3+2\epsilon+\tilde{\eta})\dot{u}_k + \frac{c_\pi^2\,k^2}{a^2}u_k = 0\,, \quad \ddot{v}_k + H(3+2\,\delta_\alpha)\dot{v}_k + \Big(m_\sigma^2 + \frac{c_\sigma^2\,k^2}{a^2}\Big)v_k = 0 \quad \text{with} \quad \delta_\alpha = \frac{\dot{\alpha}_\sigma}{H\alpha_\sigma}\,. \tag{86}$$

Their normalization follows from

$$2M_{\rm Pl}^2 H^2 \tilde{\epsilon}\, a^3\,(u_k\,\dot{u}_k^* - \dot{u}_k\,u_k^*) = i\,, \quad \alpha_\sigma^2\,a^3\,(v_k\,\dot{v}_k^* - \dot{v}_k\,v_k^*) = i\,. \tag{87}$$

Using these expressions in the interaction picture, the expectation value of $\pi_{\mathbf{k}}(t)\pi_{\mathbf{k}'}(t)$ is given by

$$\langle \pi_{\mathbf{k}}(t)\pi_{\mathbf{k}'}(t)\rangle = \langle 0|\left[\bar{T}\exp\Big(i\int_{t_0}^t dt' H_I(t')\Big)\right]\pi_{\mathbf{k}}(t)\pi_{\mathbf{k}'}(t)\left[T\exp\Big(-i\int_{t_0}^t dt' H_I(t')\Big)\right]|0\rangle \tag{88}$$

$$\begin{aligned} &= \langle 0|\pi_{\mathbf{k}}(t)\pi_{\mathbf{k}'}(t)|0\rangle + \int_{t_0}^t d\tilde{t}_1 \int_{t_0}^t dt_1 \langle 0|H^{(2)}_{\rm mix}(\tilde{t}_1)\pi_{\mathbf{k}}(t)\pi_{\mathbf{k}'}(t)H^{(2)}_{\rm mix}(t_1)|0\rangle \\ &\quad - 2{\rm Re}\left[\int_{t_0}^t dt_1 \int_{t_0}^{t_1} dt_2 \langle 0|\pi_{\mathbf{k}}(t)\pi_{\mathbf{k}'}(t)H^{(2)}_{\rm mix}(t_1)H^{(2)}_{\rm mix}(t_2)|0\rangle\right] + \dots\,, \end{aligned} \tag{89}$$

where the dots stand for the higher order terms in the couplings and $H^{(2)}_{\rm mix}$ is the second order interaction terms:

$$H^{(2)}_{\rm mix}(t) = -\int \frac{d^3\mathbf{k}}{(2\pi)^3}a^3(t)\Big[\tilde{\beta}_1\dot{\pi}_{\mathbf{k}}\,\sigma_{-\mathbf{k}} + \tilde{\beta}_2\dot{\pi}_{\mathbf{k}}\,\dot{\sigma}_{-\mathbf{k}} + \tilde{\beta}_3 c_\pi^2\frac{k^2}{a^2}\pi_{\mathbf{k}}\,\sigma_{-\mathbf{k}}\Big](t)\,. \tag{90}$$

In terms of the mode functions and the couplings, the general form of the two point function (89) is given up to the leading order corrections of the mixing couplings by

$$\langle \pi_{\mathbf{k}}(t)\pi_{\mathbf{k}'}(t)\rangle = (2\pi)^3\delta^{(3)}(\mathbf{k}+\mathbf{k}')u_k^*(t)u_k(t)\Big[1+\mathcal{C}_1+\mathcal{C}_2\Big]\,, \tag{91}$$

where $\mathcal{C}_1$ and $\mathcal{C}_2$ are defined by

$$\mathcal{C}_1 = 2\left|\int_{t_0}^t dt_1\,a^3\Big[\tilde{\beta}_1\,\dot{u}_k v_k + \tilde{\beta}_2\,\dot{u}_k\dot{v}_k + \tilde{\beta}_3 c_\pi^2\frac{k^2}{a^2}\,u_k v_k\Big](t_1)\right|^2\,, \tag{92}$$

$$\begin{aligned} \mathcal{C}_2 = -4{\rm Re}\Bigg[&\frac{u_k^2(t)}{|u_k(t)|^2}\int_{t_0}^t dt_1\,a^3\Big[\tilde{\beta}_1\,\dot{u}_k^* v_k + \tilde{\beta}_2\,\dot{u}_k^*\dot{v}_k + \tilde{\beta}_3 c_\pi^2\frac{k^2}{a^2}\,u_k^* v_k\Big](t_1) \\ &\times \int_{t_0}^{t_1} dt_2\,a^3\Big[\tilde{\beta}_1\,\dot{u}_k^* v_k^* + \tilde{\beta}_2\,\dot{u}_k^*\dot{v}_k^* + \tilde{\beta}_3 c_\pi^2\frac{k^2}{a^2}\,u_k^* v_k^*\Big](t_2)\Bigg]\,. \end{aligned} \tag{93}$$

It is convenient to rewrite $\mathcal{C} = \mathcal{C}_1 + \mathcal{C}_2$ as follows:

$$\begin{aligned} \mathcal{C} = 4{\rm Re}\Bigg[&\int_{t_0}^t dt_1\,a^3\,\Bigg[\Big(\tilde{\beta}_1\,\dot{u}_k v_k + \tilde{\beta}_2\,\dot{u}_k\dot{v}_k + \tilde{\beta}_3 c_\pi^2\frac{k^2}{a^2}\,u_k v_k\Big) - \frac{u_k^2(t)}{|u_k(t)|^2}\Big(\tilde{\beta}_1\,\dot{u}_k^* v_k + \tilde{\beta}_2\,\dot{u}_k^*\dot{v}_k + \tilde{\beta}_3 c_\pi^2\frac{k^2}{a^2}\,u_k^* v_k\Big)\Bigg](t_1) \\ &\times \int_{t_0}^{t_1} dt_2\,a^3\Big[\tilde{\beta}_1\,\dot{u}_k^* v_k^* + \tilde{\beta}_2\,\dot{u}_k^*\dot{v}_k^* + \tilde{\beta}_3 c_\pi^2\frac{k^2}{a^2}\,u_k^* v_k^*\Big](t_2)\Bigg]\,. \end{aligned} \tag{94}$$

Since the scalar perturbation ζ is given at the linear order by $\zeta = -H\pi$, we obtain the expectation value of $\zeta_{\mathbf{k}}(t)\zeta_{\mathbf{k}'}(t)$ as

$$\langle \zeta_{\mathbf{k}}(t)\zeta_{\mathbf{k}'}(t)\rangle = (2\pi)^3\delta^{(3)}(\mathbf{k}+\mathbf{k}')\frac{2\pi^2}{k^3}\mathcal{P}_\zeta(k)\,, \tag{95}$$

where the power spectrum $\mathcal{P}_\zeta(k)$ is given by

$$\mathcal{P}_\zeta(k) = \frac{H^2(t)k^3}{2\pi^2}u^*_{\mathbf{k}}(t)u_{\mathbf{k}}(t)\,(1+\mathcal{C})\,. \tag{96}$$

The factor $\mathcal{C}$ can be considered as a deviation factor from single field inflation. Here it should be noticed that in the derivation of the above general expression we assumed only $\epsilon \ll 1$, $\tilde{\epsilon} \ll 1$, and the perturbativity of the mixing couplings. In principle, we can calculate the power spectrum for any models satisfying those three conditions. In the rest of this section, we first perform the calculation concretely for the case when the time-dependence of mixing couplings is irrelevant (constant turning trajectory), and then we discuss the qualitative features of the case when the mixing couplings are large in a sufficiently short period of time (sharp turning trajectory).

A. Constant turning trajectory

In this subsection, the power spectrum is calculated in the case that the time-dependence of mixing couplings is irrelevant. To make the calculation tractable, we assume that the time-dependence of ϵ, $\tilde{\epsilon}$, α_σ, c_σ, and m_σ is negligible and we use the de-Sitter approximation. In this approximation, the equations of motion (86) for the mode functions u_k and v_k can be written as

$$u_k'' - \frac{2}{\tau}u_k' + c_\pi^2\, k^2 u_k = 0\,, \qquad v_k'' - \frac{2}{\tau}v_k' + c_\sigma^2\, k^2 v_k + \frac{m_\sigma^2}{H^2\tau^2}v_k = 0\,, \tag{97}$$

where the conformal time $d\tau = a^{-1}dt$ is given by $\tau = -1/(aH)$ in the de-Sitter approximation and the primes denote derivatives with respect to τ. The equations (97) can be solved as follows:

$$u_k = \frac{1}{2M_{\rm Pl}\tilde{\epsilon}^{1/2}(c_\pi k)^{3/2}}(1+ic_\pi k\tau)e^{-ic_\pi k\tau} = \frac{1}{2M_{\rm Pl}\tilde{\epsilon}^{1/2}(c_\pi k)^{3/2}}(1-ix)e^{ix}\,, \tag{98}$$

$$v_k = -ie^{\frac{i}{2}\pi\nu+\frac{i}{4}\pi}\frac{\sqrt{\pi}H}{2\alpha_\sigma}(-\tau)^{3/2}H_\nu^{(1)}(-c_\sigma k\tau) = -ie^{\frac{i}{2}\pi\nu+\frac{i}{4}\pi}\frac{\sqrt{\pi}H}{2\alpha_\sigma(c_\pi k)^{3/2}}x^{3/2}H_\nu^{(1)}(r_s x)\,, \tag{99}$$

where $x = -c_\pi k\tau$, $r_s = c_\sigma/c_\pi$, and we chose the Bunch-Davies vacuum for π and σ. The function $H_\nu^{(1)} = J_\nu + iY_\nu$ is the Hankel function and ν is defined as

$$\nu = \sqrt{\frac{9}{4} - \frac{m_\sigma^2}{H^2}} \quad \text{for} \quad m_\sigma < \frac{3}{2}H\,, \qquad \nu = i\sqrt{\frac{m_\sigma^2}{H^2} - \frac{9}{4}} \quad \text{for} \quad m_\sigma > \frac{3}{2}H\,. \tag{100}$$

The time derivatives of u_k and v_k are given by

$$\dot{u}_k = -H\tau u_k' = -\frac{H}{2M_{\rm Pl}\tilde{\epsilon}^{1/2}(c_\pi k)^{3/2}}x^2 e^{ix}\,, \tag{101}$$

$$\dot{v}_k = -H\tau v_k' = ie^{\frac{i}{2}\pi\nu+\frac{i}{4}\pi}\frac{\sqrt{\pi}H^2}{2\alpha_\sigma(c_\pi k)^{3/2}}x^{3/2}\Big((3/2-\nu)H_\nu^{(1)}(r_s x) + (r_s x)H_{\nu-1}^{(1)}(r_s x)\Big)\,, \tag{102}$$

where we used the identity $z\partial_z H_\nu^{(1)} = zH_{\nu-1}^{(1)} - \nu H_\nu^{(1)}$. Therefore, the factor $\mathcal{C}$ defined in (94) takes the form

$$\begin{aligned}\mathcal{C} = e^{\frac{i}{2}\pi(\nu-\nu^*)}&\frac{\pi}{4\alpha_\sigma^2}\frac{c_\pi^2}{M_{\rm Pl}^2(-\dot{H})}\\ &\times \mathrm{Re}\bigg[\int_0^\infty dx_1\Big[\Big(\frac{\tilde{\beta}_1}{H} + (\nu-3/2)^*\tilde{\beta}_2 - \tilde{\beta}_3\Big)\Big(A_1(x)^* - \widetilde{A}_1(x_1)\Big) - \tilde{\beta}_2\Big(A_2(x)^* - \widetilde{A}_2(x_1)\Big) + \tilde{\beta}_3\Big(A_3(x)^* - \widetilde{A}_3(x_1)\Big)\Big]\\ &\qquad\qquad \times \int_{x_1}^\infty dx_2\Big[\Big(\frac{\tilde{\beta}_1}{H} + (\nu-3/2)\tilde{\beta}_2 - \tilde{\beta}_3\Big)A_1(x_2) - \tilde{\beta}_2 A_2(x_2) + \tilde{\beta}_3 A_3(x_2)\Big]\bigg]\,,\end{aligned} \tag{103}$$

where we set $t_0 = -\infty$ and $t = \infty$, and we defined

$$\begin{aligned}&A_1(x) = x^{-1/2}e^{ix}H_\nu^{(1)}(r_s x)\,, \quad A_2(x) = r_s x^{1/2}e^{ix}H_{\nu-1}^{(1)}(r_s x)\,, \quad A_3(x) = ix^{1/2}e^{ix}H_\nu^{(1)}(r_s x)\,,\\ &\widetilde{A}_1(x) = x^{-1/2}e^{ix}H_{\nu^*}^{(2)}(r_s x)\,, \quad \widetilde{A}_2(x) = r_s x^{1/2}e^{ix}H_{\nu^*-1}^{(2)}(r_s x)\,, \quad \widetilde{A}_3(x) = ix^{1/2}e^{ix}H_{\nu^*}^{(2)}(r_s x)\,.\end{aligned} \tag{104}$$

When the time-dependence of mixing couplings is irrelevant, mixing couplings $\tilde{\beta}$'s can be evaluated at the time of horizon-crossing $\tau = -(c_\pi k)^{-1}$. In such a case, $\mathcal{C}$ is given by

$$\begin{aligned}\mathcal{C} = \frac{c_\pi^2}{\alpha_\sigma^2 M_{\rm Pl}^2(-\dot{H})}{\rm Re}\Bigg[&\left|\tilde{\beta}_1/H + (\nu - 3/2)\tilde{\beta}_2 - \tilde{\beta}_3\right|^2 \mathcal{I}_{11} + (\tilde{\beta}_1/H + (\nu^* - 3/2)\tilde{\beta}_2 - \tilde{\beta}_3)(-\tilde{\beta}_2\,\mathcal{I}_{12} + \tilde{\beta}_3\,\mathcal{I}_{13}) \\ &- \tilde{\beta}_2\Big((\tilde{\beta}_1/H + (\nu - 3/2)\tilde{\beta}_2 - \tilde{\beta}_3)\mathcal{I}_{21} - \tilde{\beta}_2\,\mathcal{I}_{22} + \tilde{\beta}_3\,\mathcal{I}_{23}\Big) \\ &+ \tilde{\beta}_3\Big((\tilde{\beta}_1/H + (\nu - 3/2)\tilde{\beta}_2 - \tilde{\beta}_3)\mathcal{I}_{31} - \tilde{\beta}_2\,\mathcal{I}_{32} + \tilde{\beta}_3\,\mathcal{I}_{33}\Big)\Bigg] \qquad (105)\end{aligned}$$

$$= \frac{c_\pi^2}{\alpha_\sigma^2 M_{\rm Pl}^2(-\dot{H})}\left(\frac{\tilde{\beta}_1^2}{H^2}\,\mathcal{C}_{11} + \tilde{\beta}_2^2\,\mathcal{C}_{22} + \tilde{\beta}_3^2\,\mathcal{C}_{33} + \frac{\tilde{\beta}_1}{H}\,\tilde{\beta}_2\,\mathcal{C}_{12} + \frac{\tilde{\beta}_1}{H}\,\tilde{\beta}_3\,\mathcal{C}_{13} + \tilde{\beta}_2\,\tilde{\beta}_3\,\mathcal{C}_{23}\right), \qquad (106)$$

where $\mathcal{I}_{ij}$ is integral defined as

$$\mathcal{I}_{ij} = \frac{\pi}{4}e^{\frac{i}{2}\pi(\nu-\nu^*)}\int_0^\infty dx_1\Big(A_i(x_1)^* - \widetilde{A}_i(x_1)\Big)\int_{x_1}^\infty dx_2\,A_j(x_2)\,, \qquad (107)$$

and $\mathcal{C}_{ij}$ is given by

$$\begin{aligned}&\mathcal{C}_{11} = {\rm Re}\Big[\,\mathcal{I}_{11}\,\Big]\,,\quad \mathcal{C}_{22} = {\rm Re}\Big[|\nu - 3/2|^2\,\mathcal{I}_{11} - (\nu^* - 3/2)\,\mathcal{I}_{12} - (\nu - 3/2)\,\mathcal{I}_{21} + \mathcal{I}_{22}\Big]\,, \\ &\mathcal{C}_{33} = {\rm Re}\left[\,\mathcal{I}_{11} + \mathcal{I}_{33} - \mathcal{I}_{13} - \mathcal{I}_{31}\,\right]\,,\quad \mathcal{C}_{12} = {\rm Re}\left[(\nu + \nu^* - 3)\,\mathcal{I}_{11} - \mathcal{I}_{12} - \mathcal{I}_{21}\right]\,,\quad \mathcal{C}_{13} = {\rm Re}\left[-2\,\mathcal{I}_{11} + \mathcal{I}_{13} + \mathcal{I}_{31}\right]\,, \\ &\mathcal{C}_{23} = {\rm Re}\left[-(\nu + \nu^* - 3)\,\mathcal{I}_{11} + \mathcal{I}_{12} + \mathcal{I}_{21} + (\nu^* - 3/2)\,\mathcal{I}_{13} + (\nu - 3/2)\,\mathcal{I}_{31} - \mathcal{I}_{23} - \mathcal{I}_{32}\right]\,. \qquad (108)\end{aligned}$$

Here the explicit form of the power spectrum is given by

$$\mathcal{P}_\zeta(k) = \frac{H^2}{8\pi^2 M_{\rm Pl}^2 \epsilon\, c_\pi}\left[1 + \frac{c_\pi^2}{\alpha_\sigma^2 M_{\rm Pl}^2(-\dot{H})}\left(\frac{\tilde{\beta}_1^2}{H^2}\,\mathcal{C}_{11} + \tilde{\beta}_2^2\,\mathcal{C}_{22} + \tilde{\beta}_3^2\,\mathcal{C}_{33} + \frac{\tilde{\beta}_1}{H}\,\tilde{\beta}_2\,\mathcal{C}_{12} + \frac{\tilde{\beta}_1}{H}\,\tilde{\beta}_3\,\mathcal{C}_{13} + \tilde{\beta}_2\,\tilde{\beta}_3\,\mathcal{C}_{23}\right)\right], \qquad (109)$$

where H, ϵ, c_π, α_σ, $\tilde{\beta}_i$, and r_s are evaluated at the time of horizon-crossing. Then, the calculation of the power spectrum reduces to the evaluation of ${\rm Re}[\mathcal{I}_{ij} + \mathcal{I}_{ji}]$, ${\rm Im}[\mathcal{I}_{12} - \mathcal{I}_{21}]$, and ${\rm Im}[\mathcal{I}_{13} - \mathcal{I}_{31}]$.

As is understood from the definition, $\mathcal{I}_{ij}$'s and $\mathcal{C}_{ij}$'s are functions of m_σ and $r_s = c_\sigma/c_\pi$. For general value of r_s, it is difficult to perform the integrals analytically and we performed numerical calculations by contour deformations (see appendix A for details). For the special case $r_s = 1$, however, it is possible to perform the integrals $\mathcal{I}_{ij}$'s analytically by extending the results in [22], and the results are summarized in appendix B. In such a way, $\mathcal{C}_{ij}$'s are calculated and the obtained results are summarized in figure 1 and figure 2. We find that they monotonically decrease in m_σ for fixed r_s, but they are not monotonic for fixed m_σ. It is also notable that $\mathcal{C}_{22}$ does not vanish even in the limit $m_\sigma \to \infty$ contrary to the usual expectation that heavy particles decouple for finite mixing couplings. In the last of this subsection, we give an explanation of the peculiar features of heavy particles.

In the heavy mass limit $m_\sigma \gg H$ and $m_\sigma \gg a^{-1}c_\sigma k$, the mode function v_k can be expressed as

$$v_k \sim \frac{1}{\alpha_\sigma\sqrt{2m_\sigma}}a^{-3/2}e^{-im_\sigma t}\,, \qquad (110)$$

where the normalization of v_k follows from (87). Then, the mixing couplings $\tilde{\beta}_1$ and $\tilde{\beta}_3$ are suppressed by the prefactor $1/\sqrt{m_\sigma}$ and these mixings do not contribute to the deviation factor $\mathcal{C}$ in the heavy mass limit. On the other hand, in the case of the $\tilde{\beta}_2$ mixing coupling, such a suppression does not occur because $\dot{v}_k$ takes the form[8]

$$\dot{v}_k \sim -i\frac{\sqrt{m_\sigma}}{\sqrt{2}\alpha_\sigma}a^{-3/2}e^{-im_\sigma t}\,. \qquad (111)$$

---

[8] Apparently, the expression (111) seems to imply a divergence in the mixing. However, it turns out that such a divergence does not occur because of the integration over time.



FIG. 1: $\mathcal{C}_{ij}$'s for fixed $r_s = c_\sigma/c_\pi$. The dots are numerical results for $r_s = 0.1$ (red), 0.3 (orange), 1 (yellow), 3 (green), and 10 (blue). The curves are analytic results for $r_s = 1$ obtained in the next subsection.

More explicitly, it is possible to evaluate $\mathcal{C}_{22}$ in the heavy mass limit. In this limit, v_k and $\dot{v}_k$ are related to each other as

$$\dot{v}_k = -im_\sigma v_k\,, \tag{112}$$

and $\mathcal{C}_{22}$ is related to $\mathcal{C}_{11}$ as

$$\mathcal{C}_{22} = \frac{m_\sigma^2}{H^2}\mathcal{C}_{11}\,. \tag{113}$$

As was discussed in [18, 22], $\mathcal{C}_{11}$ for $r_s = 1$ in the heavy mass limit is analytically evaluated as[9]

$$\mathcal{C}_{11} = \frac{1}{4}\frac{H^2}{m_\sigma^2}\,. \tag{114}$$

We then obtain

$$\mathcal{C}_{22} = \frac{1}{4} \quad \text{in the heavy mass limit.} \tag{115}$$

---

[9] The factor $\mathcal{C}(\mu)$ in [18, 22] corresponds to our $\mathcal{C}_{11}$ for $r_s = 1$ and $\mu \sim \dfrac{m_\sigma}{H}$ in the heavy mass limit.



FIG. 2: $\mathcal{C}_{ij}$'s for fixed m_σ. The dots are numerical results for $m_\sigma/H = 1.2$ (red), 1.6 (yellow), 2.0 (green), and 4.0 (blue).

Note that, in the heavy mass limit, v_k and therefore $\mathcal{C}_{ij}$'s do not depend on r_s. The result (115) is consistent with our numerical and analytic results and we conclude that the $\tilde{\beta}_2$ mixing coupling contributes to the deviation factor $\mathcal{C}$ even in the heavy mass limit. It would be important to understand this feature more physically.

B. Qualitative features of sharp turning trajectory

In this subsection, we discuss qualitative features of the case when the mixing couplings are large in a sufficiently short period of time (sharp turning trajectory). As a simplest example, let us first consider the case when the $\tilde{\beta}_1$ coupling is the only relevant mixing coupling and it is proportional to a delta function:

$$\tilde{\beta}_1 = \beta_* \,\delta(t - t_*) = \beta_* a^{-1}(t_*)\,\delta(\tau - \tau_*)\,, \quad \tilde{\beta}_2 = \tilde{\beta}_3 = 0\,, \tag{116}$$

where t_* is the time of sharp turning, τ_* is the corresponding conformal time, and the mode k_* crossing the horizon at $t = t_*$ is given by $k_* = -1/(c_\pi \tau_*)$. We assume that β_* is small enough to be treated perturbatively. In this case, the deviation of the power spectrum from that of single field inflation can be written as[10]

$$\mathcal{C} = 2\beta_*^2 \,\mathrm{Re}\left[a^6\left(\dot{u}_k^* - \dot{u}_k\right)\dot{u}_k\, v_k\, v_k^*(t_*)\right], \tag{117}$$

which is related to the power spectrum by (96). For simplicity, suppose that time-dependence of m_σ, α_σ, c_σ, and c_π is negligible at the time of sharp turning and the mode functions $u_k(t_*)$ and $v_k(t_*)$ are given by

$$u_k(t_*) = \frac{1}{2M_{\mathrm{Pl}}\tilde{\epsilon}^{1/2}(c_\pi k)^{3/2}}(1 - ix_*)e^{ix_*}\,, \tag{118}$$

$$v_k(t_*) = -ie^{\frac{i}{2}\pi\nu + \frac{i}{4}\pi}\frac{\sqrt{\pi}H}{2\alpha_\sigma(c_\pi k)^{3/2}}x_*^{3/2}H_\nu^{(1)}(r_s x_*)\,, \tag{119}$$

where $x_* = k/k_*$ and parameters such as m_σ are evaluated at the time of sharp turning. Then, (117) is given by

$$\mathcal{C} = \beta_*^2\,\frac{\pi}{4\alpha_\sigma^2}\frac{c_\pi^2}{M_{\mathrm{Pl}}^2(-\dot{H})}\mathcal{F}_\nu(x_*) \quad \text{with} \quad \mathcal{F}_\nu(x_*) = e^{\frac{i}{2}\pi(\nu-\nu^*)}x_*\sin^2 x_*|H_\nu^{(1)}(r_s x_*)|^2\,. \tag{120}$$

For $k \ll k_*$ or $x_* \ll 1$, the asymptotic behavior of $\mathcal{F}_\nu(x_*)$ is

$$\mathcal{F}_\nu(x_*) \sim \begin{cases} \dfrac{4^\nu}{\pi^2}\Gamma(\nu)^2\, r_s^{-2\nu}x_*^{3-2\nu} & \text{for} \quad m_\sigma(t_*) < \dfrac{3}{2}H\,, \\[2ex] \dfrac{1}{\pi^2}\left|e^{\frac{i}{2}\pi\nu}2^\nu\Gamma(\nu)(r_s x_*)^{-\nu} + e^{-\frac{i}{2}\pi\nu}2^{-\nu}\Gamma(-\nu)(r_s x_*)^\nu\right|^2 x_*^3 & \text{for} \quad m_\sigma(t_*) > \dfrac{3}{2}H\,, \end{cases} \tag{121}$$

which is consistent to the intuition that modes outside the horizon at $t = t_*$ are not much affected by the sharp turning. For $k \gg k_*$ or $x_* \gg 1$, it reduces to

$$\mathcal{F}_\nu(x_*) \sim \frac{2}{\pi}r_s^{-1}\sin^2 x_* = \frac{1}{\pi}r_s^{-1}(1 - \cos 2x_*)\,. \tag{122}$$

This kind of oscillating behavior was also found in [15–17, 19]. The turning trajectory generically oscillates around the turning point and the mixing couplings at the turning point become more regular than delta functions. In such a case, it is expected that the oscillating behavior of short modes $c_\pi k \gtrsim -1/\tau_*$ begins damping at some scale characterized by the oscillation of the trajectory. Let us next confirm such a behavior explicitly for a concrete example with a finite width of turning. We consider the following $\tilde{\beta}_1$ profile:

$$\tilde{\beta}_1 = \begin{cases} \beta_*(-H\tau_*)(\Delta\tau)^{-1} & \text{for} \quad \tau_* - \frac{\Delta\tau}{2} < \tau < \tau_* + \frac{\Delta\tau}{2}\,, \\ 0 & \text{otherwise}\,, \end{cases} \tag{123}$$

where we normalized β_* so that it reproduces the coupling (116) in the limit $\Delta\tau \to 0$. For this class of couplings, we define $\mathcal{F}_\nu$ by

$$\mathcal{C} = \beta_*^2\,\frac{\pi}{4\alpha_\sigma^2}\frac{c_\pi^2}{M_{\mathrm{Pl}}^2(-\dot{H})}\mathcal{F}_\nu(x_*)\,. \tag{124}$$

Since it is difficult to calculate $\mathcal{F}_\nu$ analytically, we can confirm the expected damping behavior of $\mathcal{F}_\nu$ by numerical calculations and the results are given in figure 3. To summarize, $\mathcal{C}$ vanishes in the long mode limit $c_\pi k \ll -1/\tau_*$, oscillates for short modes $c_\pi k \gtrsim -1/\tau_*$, and damps at some scale characterized by the oscillating trajectory around the turning point.

It is straightforward to extend the above discussions to the case with non-vanishing $\tilde{\beta}_2$ and $\tilde{\beta}_3$. We generically expect to find a similar behavior of $\mathcal{C}$: vanishing in the long mode limit $c_\pi k \ll -1/\tau_*$, oscillating for short modes $c_\pi k \gtrsim -1/\tau_*$, and damping at some scale characterized by the oscillating trajectory around the turning point.

---

[10] In our calculation we have treated the mixing coupling β_* as an interaction. In such an interaction picture, it is manifest that the deviation from single field inflation starts from the second order in β_*. On the other hand, it is also possible to treat the mixing coupling as a part of the kinetic and mass terms. In that picture, the commutation relations (85) of creation and annihilation operators are affected by the mixing as well as the mode functions u_k and v_k are modified. However, in some literatures, these modifications are not taken into account adequately and the deviation from single field inflation is calculated to start from the first order in β_*.



FIG. 3: $\mathcal{F}_\nu$ for fixed $c_\pi k_* \Delta\tau$ and $c_\pi = c_\sigma$

. The left figure is for $m_\sigma = H$ and the right one is for $m_\sigma = 3H$. The dots are numerical results for $c_\pi k_* \Delta\tau = 5$ (red), 10 (green), and 15 (blue). The curves are analytic results in the limit of $c_\pi k_* \Delta \to 0$ (delta function limit).

V. THREE POINT FUNCTIONS IN THE SQUEEZED LIMIT

In this section, we discuss the momentum dependence of three point functions in the squeezed limit. We take the decoupling limit and assume that time dependence of α's, β's, γ's, and $\dot{H}$ is negligible. Under these assumption, the second order action is given in (82) and (83), and three point vertices are given by

$$\begin{aligned} S^{(3)} = \int d^4x\, a^3 \Bigg[&-\Big(M_{\rm Pl}^2 \dot{H}(c_\pi^{-2}-1) + \frac{4M_3^4}{3}\Big)\dot{\pi}^3 + M_{\rm Pl}^2\dot{H}(c_\pi^{-2}-1)\dot{\pi}\frac{(\partial_i\pi)^2}{a^2} + (-\beta_1 + 4\overline{\gamma}_6)\dot{\pi}^2\sigma + (3\beta_2 - 4\overline{\gamma}_7)\dot{\pi}^2\dot{\sigma} \\ &- 2\beta_2\dot{\pi}\frac{\partial_i\pi\partial_i\sigma}{a^2} + \beta_1\frac{(\partial_i\pi)^2}{a^2}\sigma - \beta_2\frac{(\partial_i\pi)^2}{a^2}\dot{\sigma} - 2\overline{\gamma}_1\dot{\pi}\sigma^2 + (-\alpha_4 + 2\overline{\gamma}_2)\dot{\pi}\sigma\dot{\sigma} \\ &+ \Big(\alpha_\sigma^2(1-c_\sigma^2) - 2\overline{\gamma}_3 + 2\overline{\gamma}_5\Big)\dot{\pi}\dot{\sigma}^2 - 2\overline{\gamma}_5\dot{\pi}\frac{(\partial_i\sigma)^2}{a^2} + \alpha_4\frac{\partial_i\pi\partial_i\sigma}{a^2}\sigma - \alpha_\sigma^2(1-c_\sigma^2)\frac{\partial_i\pi\partial_i\sigma}{a^2}\dot{\sigma} - 2\overline{\gamma}_4\ddot{\pi}\sigma\dot{\sigma} \\ &+ \gamma_1\sigma^3 - \gamma_2\sigma^2\dot{\sigma} + (\gamma_3-\gamma_5)\sigma\dot{\sigma}^2 + \gamma_5\sigma\frac{(\partial_i\sigma)^2}{a^2} + (-\gamma_4+\gamma_6)\dot{\sigma}^3 - \gamma_6\dot{\sigma}\frac{(\partial_i\sigma)^2}{a^2}\Bigg]\,. \end{aligned} \tag{125}$$

In the following, we discuss what kind of momentum dependence in the squeezed limit appears from the above three point vertices.

As an example, let us start from the case when the mixing coupling $\tilde{\beta}_1\dot{\pi}\sigma$ and the three point vertex $4\bar{\gamma}_6\dot{\pi}^2\sigma$ are relevant. In this case, the three point function of π takes the form

$$\begin{aligned} &\langle \pi_{\mathbf{k}_1}(t)\pi_{\mathbf{k}_2}(t)\pi_{\mathbf{k}_3}(t)\rangle \\ &\ni \frac{\tilde{\beta}_1\,\bar{\gamma}_6\,(2\pi)^3\delta^{(3)}(\mathbf{k}_1+\mathbf{k}_2+\mathbf{k}_3)}{M_{\rm Pl}^3\tilde{\epsilon}^{3/2}(c_\pi k_1)^{3/2}(c_\pi k_2)^{3/2}(c_\pi k_3)^{3/2}}{\rm Re}\Bigg[\int_{-\infty}^{\infty} dt_1\, a^3\,\dot{u}_{k_1}^*(t_1)\dot{u}_{k_2}^*(t_1) \\ &\quad\times\Bigg\{v_{k_3}^*(t_1)\int_{-\infty}^{\infty}dt_2\,a^3\,\dot{u}_{k_3}(t_2)v_{k_3}(t_2) - v_{k_3}^*(t_1)\int_{t_1}^{\infty}dt_2\,a^3\,\dot{u}_{k_3}^*(t_2)v_{k_3}(t_2) - v_{k_3}(t_1)\int_{-\infty}^{t_1}dt_2\,a^3\,\dot{u}_{k_3}^*(t_2)v_{k_3}^*(t_2)\Bigg\}\Bigg] \\ &\quad+ (2\ \text{permutations})\,. \end{aligned} \tag{126}$$

Here there are three terms in the curly brackets, and they correspond to the Feynman diagrams in figure 4, respectively. In the squeezed limit, $k_1 = k_2 = k$ and $k_3/k = \kappa \ll 1$, the long mode k_3 crosses the horizon much earlier than the short modes k_1 and k_2. Since the interactions are considered to be relevant around the horizon, it is expected that the relevant contribution arises from $(\tau_1, \tau_2) \sim (-1/k_1, -1/k_3)$, and therefore, the middle term in the curly brackets becomes irrelevant in the squeezed limit. In fact, we can confirm this expectation explicitly from the expression (126), and the integrals in the curly brackets can be written at the leading order in κ as

$$\int_{-\infty}^{\infty} dt_1\, a^3\,\dot{u}_{k_1}^*(t_1)\dot{u}_{k_2}^*(t_1)\,2i\,{\rm Im}\left[v_{k_3}^*(t_1)\int_{-\infty}^{\infty}dt_2\,a^3\,\dot{u}_{k_3}(t_2)v_{k_3}(t_2)\right]\,. \tag{127}$$



FIG. 4: Feynman diagrams for the first term of (126). The solid lines denote the propagations of π and the dotted lines denote those of σ.

We notice that the t_2-integral is k_3-independent and k_3-dependence of the total integral appears only via $v^*_{k_3}(t_1)$ originated from the three-point vertex. Then, the momentum dependence of the first term in (126) is given by

$$\text{the first term in (126)} \;\propto \begin{cases} \kappa^{-3/2-\nu}k^{-6} & \text{for} \quad m_\sigma < \dfrac{3}{2}H\,, \\ \kappa^{-3/2}k^{-6}\sin[i\nu\log\kappa + \delta_\nu] & \text{for} \quad m_\sigma > \dfrac{3}{2}H\,, \end{cases} \tag{128}$$

where δ_ν is a ν-dependent phase factor. We note that the only information necessary to derive (128) was the fact that the field of momentum k_3 in the three point vertex takes the form σ. For the other two permutation terms in (126), the k_3-dependence of the integral is determined by $u^*_{k_3}$ originated from the three point vertex, and their momentum dependence is given by

$$\text{the other two permutation terms in (126)} \;\propto \kappa^{-1}k^{-6}\,. \tag{129}$$

Then, the first term dominates for small κ. Therefore, when the mixing coupling $\tilde{\beta}_1$ and three point coupling $\overline{\gamma}_6$ are relevant, the momentum dependence of scalar three point functions in the squeezed limit is given by

$$\lim_{k_3/k_1=k_3/k_2=\kappa\to 0}\langle\zeta_{\mathbf{k}_1}\zeta_{\mathbf{k}_2}\zeta_{\mathbf{k}_3}\rangle \propto \begin{cases} \kappa^{-3/2-\nu}k^{-6} & \text{for} \quad m_\sigma < \dfrac{3}{2}H\,, \\ \kappa^{-3/2}k^{-6}\sin[i\nu\log\kappa + \delta_\nu] & \text{for} \quad m_\sigma > \dfrac{3}{2}H\,. \end{cases} \tag{130}$$

It is straightforward to extend the above discussion for more general cases. First, for general $\tilde{\beta}_i$'s, we can show that when the mixing couplings convert π of momentum $k_3 = \kappa k$ to σ, the t_2-integral becomes κ-independent in the limit $\kappa \ll 1$ and the three point vertex determines the κ-dependence of the diagram. Second, the only information necessary to obtain the momentum dependence is the form of the field of momentum k_3 in the vertex. In the previous examples, when it takes the form σ in the vertex the diagram was proportional to $\kappa^{-3/2-\nu}$ or $\kappa^{-3/2}\sin[i\nu\log\kappa+\delta_\nu]$, and it was proportional to κ^{-1} for $\dot{\pi}$. More generally, we can obtain the relations in Table I between momentum dependence of the diagram and the form of the field of momentum k_3 in the three point vertex. Here it should be noted that the ν-dependent phase factor δ_ν depends on the details of mixing couplings. Finally, as discussed in the previous example, momentum dependence of the contribution from each vertex is now identified for $\kappa \ll 1$, it is straightforward to obtain momentum dependence of the contribution to scalar three point functions from each three point vertex displayed in (125). The results are summarized in Table II. Here note that although the contribution from the $\dot{\pi}\dfrac{(\partial_i\pi)^2}{a^2}$ vertex seems to be proportional to $\kappa^{-2}k^{-6}$ apparently, explicit calculations show that this kind of leading contribution vanishes and the three point functions begin with terms proportional to $\kappa^{-1}k^{-6}$.

As we have seen, the momentum dependence of scalar three point functions in the squeezed limit has robust information about mass of σ and three point vertices.

VI. SUMMARY AND DISCUSSION

In this paper we discussed quasi-single field inflation based on the effective field theory approach. We first constructed the most generic action in the unitary gauge based on the unbroken time-dependent spatial diffeomorphism,

form of the field of momentum k_3	momentum dependence of the diagram
π	$\kappa^{-3}k^{-6}$
$\dot{\pi}$, $\ddot{\pi}$	$\kappa^{-1}k^{-6}$
$\dfrac{\partial_i\pi}{a}$	$\kappa^{-2}k^{-6}$
σ, $\dot{\sigma}$	$\begin{cases} \kappa^{-3/2-\nu}k^{-6} & \text{for} \quad m_\sigma < \frac{3}{2}H \\ \kappa^{-3/2}k^{-6}\sin[i\nu\log\kappa+\delta_\nu] & \text{for} \quad m_\sigma > \frac{3}{2}H \end{cases}$
$\dfrac{\partial_i\sigma}{a}$	$\begin{cases} \kappa^{-1/2-\nu}k^{-6} & \text{for} \quad m_\sigma < \frac{3}{2}H \\ \kappa^{-1/2}k^{-6}\sin[i\nu\log\kappa+\delta_\nu] & \text{for} \quad m_\sigma > \frac{3}{2}H \end{cases}$

TABLE I: Momentum dependence of the diagram

three point vertices	momentum dependence
$\dot{\pi}^3$, $\dot{\pi}\dfrac{(\partial_i\pi)^2}{a^2}$	$\kappa^{-1}k^{-6}$
$\dot{\pi}^2\sigma$, $\dot{\pi}^2\dot{\sigma}$, $\dot{\pi}\sigma^2$, $\dot{\pi}\sigma\dot{\sigma}$, $\dot{\pi}\dot{\sigma}^2$, $\ddot{\pi}\sigma\dot{\sigma}$, σ^3, $\sigma^2\dot{\sigma}$, $\sigma\dot{\sigma}^2$, $\sigma\dfrac{(\partial_i\sigma)^2}{a^2}$, $\dot{\sigma}^3$, $\dot{\sigma}\dfrac{(\partial_i\sigma)^2}{a^2}$	$\begin{cases} \kappa^{-3/2-\nu}k^{-6} & \text{for} \quad m_\sigma < \frac{3}{2}H \\ \kappa^{-3/2}k^{-6}\sin[i\nu\log\kappa+\delta_\nu] & \text{for} \quad m_\sigma > \frac{3}{2}H \end{cases}$
$\dot{\pi}\dfrac{\partial_i\pi\partial_i\sigma}{a^2}$	$\kappa^{-2}k^{-6}$
$\dfrac{(\partial_i\pi)^2}{a^2}\sigma$, $\dfrac{(\partial_i\pi)^2}{a^2}\dot{\sigma}$, $\dfrac{\partial_i\pi\partial_i\sigma}{a^2}\sigma$, $\dfrac{\partial_i\pi\partial_i\sigma}{a^2}\dot{\sigma}$	$\begin{cases} \kappa^{-3/2-\nu}k^{-6} & \text{for} \quad m_\sigma < \sqrt{2}H \\ \kappa^{-2}k^{-6} & \text{for} \quad m_\sigma > \sqrt{2}H \end{cases}$
$\dot{\pi}\dfrac{(\partial_i\sigma)^2}{a^2}$	$\begin{cases} \kappa^{-1/2-\nu}k^{-6} & \text{for} \quad m_\sigma < \sqrt{2}H \\ \kappa^{-1}k^{-6} & \text{for} \quad m_\sigma > \sqrt{2}H \end{cases}$

TABLE II: Three point vertices and momentum dependence

and then constructed the action for the Goldstone boson by the Stückelberg method. Its decoupling regime was also discussed carefully, and the action in the decoupling regime implies that non-trivial cubic interactions generically appear and large non-Gaussianities can arise when the sound speed c_σ of σ is small, α_4 is non-zero, or mixing couplings β_1 and β_2 exist as well as the sound speed c_π of π is small. Using the obtained action, two classes of concrete models were discussed: the constant turning trajectory and the sharp turning trajectory.

In the constant turning case, we first calculated the power spectrum of scalar perturbations numerically for general values of $r_s = c_\sigma/c_\pi$ and analytically for the special case $r_s = 1$. It was found that when the $\dot{\pi}\dot{\sigma}$ coupling is relevant, the massive scalar field σ affects the dynamics of the Goldstone π even in the heavy mass limit $m_\sigma \gg H$ contrary to the usual expectation that heavy particles decouple for finite mixing couplings. We also discussed the momentum dependence of scalar three point functions in the squeezed limit for general settings of quasi-single field inflation. It was shown that the momentum dependence is determined only from the cubic interactions and the cubic interactions were classified into five classes. The three point functions in the squeezed limit take the intermediate shapes between local and equilateral types when the mixing couplings are relevant, and this kind of momentum dependence characterizes quasi-single field inflation. Recently in [32], the detectability of such a momentum dependence was discussed for some cases. It would be interesting to discuss the detectability of the momentum dependence in the form of $\kappa^{-3/2}\sin[i\nu\log\kappa+\delta_\nu]$, which arises in the second class with $m_\sigma > \frac{3}{2}H$. It is also important to calculate the full bi-spectrum for general settings of quasi-single field inflation.

In the sharp turning case, we made a qualitative discussion of the power spectrum. It was found that the deviation factor $\mathcal{C}$ from single field inflation vanishes for long modes $c_\pi k \ll -1/\tau_*$, oscillates for short modes $c_\pi k \gtrsim -1/\tau_*$, and damps at some scale characterized by the oscillating trajectory around the turning point. Since our framework makes the contributions from the mixing couplings clear, it would be useful to discuss more on the sharp turning trajectory.

Our framework can be considered as a starting point for systematic discussions on multiple field models, and there would be a lot of applications such as those mentioned above. We hope to report our progress in these directions elsewhere.

Acknowledgments

The work of T.N. was supported in part by JSPS Grant-in-Aid for JSPS Fellows. This work was supported in part by the Grant-in-Aid for Scientific Research No. 21740187 (M.Y.) and the Grant-in-Aid for Scientific Research on Innovative Areas No. 24111706 (M.Y.). D.Y. acknowledges the financial support from the Global Center of Excellence Program by MEXT, Japan through the "Nanoscience and Quantum Physics" Project of the Tokyo Institute of Technology.

Appendix A: Numerical calculations of power spectrum

To perform the integral (107) numerically, there are two technical obstacles [21, 22]: spurious divergences at $x = 0$ and oscillating behaviors at $x = \infty$.

The integral (107) contains two integrals, $\int_0^\infty dx_1\, A_i(x_1)^* \int_{x_1}^\infty dx_2\, A_j(x_2)$ and $\int_0^\infty dx_1\, \widetilde{A}_i(x_1) \int_{x_1}^\infty dx_2\, A_j(x_2)$, and each integral diverges for some parameter region. However, we can show that such divergences cancel out in the calculation of $\mathcal{C}_{ij}$'s. For example, let us consider $\mathcal{C}_{11}$ for real $0 < \nu < 3/2$. The asymptotic behavior of each integral in $\mathcal{I}_{11}$ around $x_1 = x_2 = 0$ is given by

$$\sim \int_0 dx_1\, x_1^{-1/2-\nu} \int_{x_1} dx_2\, x_2^{-1/2-\nu} \sim 0^{1-2\nu}\,, \tag{A1}$$

which diverges for $\nu > 1/2$. We first notice that this kind of leading singularities cancel out between two terms and the asymptotic behavior of the total integral $\mathcal{I}_{11}$ is given by

$$\sim i \int_0 dx_1\, x_1^{1/2-\nu} \int_{x_1} dx_2\, x_2^{-1/2-\nu} \sim i\, 0^{2-2\nu} \tag{A2}$$

up to some real constant number. Although it is still singular for $\nu > 1$, such singular contribution is pure-imaginary and does not contribute to $\mathrm{Re}[\mathcal{I}_{11}]$. The higher order terms are finite for $\nu < 3/2$ and hence we conclude that $\mathcal{C}_{11} = \mathrm{Re}[\mathcal{I}_{11}]$ is finite. In a similar way, we can show that $\mathrm{Re}[\mathcal{I}_{\mathrm{ij}} + \mathcal{I}_{\mathrm{ji}}]$, $\mathrm{Im}[\mathcal{I}_{12} - \mathcal{I}_{21}]$, and $\mathrm{Im}[\mathcal{I}_{13} - \mathcal{I}_{31}]$ are finite for $0 < \nu < 3/2$ and $\nu =$ pure-imaginary, and therefore all the $\mathcal{C}_{ij}$'s are finite. To avoid this kind of spurious singularities in the numerical calculation, we introduce a IR cut off ϵ_{IR}:

$$\mathcal{I}_{ij} = \frac{\pi}{4} e^{\frac{i}{2}\pi(\nu-\nu^*)} \int_{\epsilon_{\mathrm{IR}}}^\infty dx_1 \Big(A_i(x_1)^* - \widetilde{A}_i(x_1) \Big) \int_{x_1}^\infty dx_2\, A_j(x_2)\,, \tag{A3}$$

where we set $\epsilon_{\mathrm{IR}} = 10^{-10}$ in our calculation.

As is usual in the Feynman diagram calculation in the momentum space, the integral $\mathcal{I}_{ij}$ oscillates at $x = \infty$ because of the oscillating behavior of the mode functions u_k and v_k, which makes the numerical calculation difficult. Following [21, 22], we perform contour deformations to avoid this kind of technical difficulties. Let us first consider the integral $\mathrm{Re}[\mathcal{I}_{ij}+\mathcal{I}_{ji}]$. It is convenient to rewrite it as follows:

$$\begin{aligned}\mathrm{Re}[\mathcal{I}_{ij}+\mathcal{I}_{ji}] &= \frac{\pi}{4}e^{\frac{i}{2}\pi(\nu-\nu^*)}\mathrm{Re}\bigg[\int_{\epsilon_{\mathrm{IR}}}^{\infty}dx_1\,A_i(x_1)^*\int_{x_1}^{\infty}dx_2\,A_j(x_2)+\int_{\epsilon_{\mathrm{IR}}}^{\infty}dx_1\,A_j(x_1)\int_{x_1}^{\infty}dx_2\,A_i(x_2)^* \\ &\qquad -\int_{\epsilon_{\mathrm{IR}}}^{\infty}dx_1\,\widetilde{A}_i(x_1)\int_{x_1}^{\infty}dx_2\,A_j(x_2)-\int_{\epsilon_{\mathrm{IR}}}^{\infty}dx_1\,\widetilde{A}_j(x_1)\int_{x_1}^{\infty}dx_2\,A_i(x_2)\bigg] \\ &= \frac{\pi}{4}e^{\frac{i}{2}\pi(\nu-\nu^*)}\mathrm{Re}\bigg[\bigg(\int_{\epsilon_{\mathrm{IR}}}^{\infty}dx\,A_i(x)\bigg)^*\int_{\epsilon_{\mathrm{IR}}}^{\infty}dy\,A_j(y) \\ &\qquad -\int_{\epsilon_{\mathrm{IR}}}^{\infty}dx_1\,\widetilde{A}_i(x_1)\int_{x_1}^{\infty}dx_2\,A_j(x_2)-\int_{\epsilon_{\mathrm{IR}}}^{\infty}dx_1\,\widetilde{A}_j(x_1)\int_{x_1}^{\infty}dx_2\,A_i(x_2)\bigg]\,. \end{aligned}\tag{A4}$$

The first term in the bracket can be Wick rotated without crossing any poles as

$$\bigg(\int_0^{\infty}dx\,A_i(\epsilon_{\mathrm{IR}}+ix)\bigg)^*\int_0^{\infty}dy\,A_j(\epsilon_{\mathrm{IR}}+iy)\,,\tag{A5}$$

and the last two terms are Wick rotated as

$$\int_0^{\infty}dx\,\widetilde{A}_i(\epsilon_{\mathrm{IR}}+ix)\int_0^{\infty}dy\,A_j(\epsilon_{\mathrm{IR}}+ix+iy)+\int_0^{\infty}dx\,\widetilde{A}_j(\epsilon_{\mathrm{IR}}+ix)\int_0^{\infty}dy\,A_i(\epsilon_{\mathrm{IR}}+ix+iy)\,.\tag{A6}$$

Then, we obtain the following expression of $\mathrm{Re}[\mathcal{I}_{ij}+\mathcal{I}_{ji}]$:

$$\mathrm{Re}[\mathcal{I}_{ij}+\mathcal{I}_{ji}]=\frac{\pi}{4}e^{\frac{i}{2}\pi(\nu-\nu^*)}\mathrm{Re}\bigg[\int_0^{\infty}dx\Big(A_i(\epsilon_{\mathrm{IR}}+ix)^*+\widetilde{A}_i(\epsilon_{\mathrm{IR}}+ix)\Big)\int_0^{\infty}dy\,A_j(\epsilon_{\mathrm{IR}}+ix+iy)\bigg]+(i\leftrightarrow j)\,.\tag{A7}$$

To avoid the singular behavior around $x=0$ discussed above, we further modify the contour as follows:

$$\begin{aligned}\mathrm{Re}[\mathcal{I}_{ij}+\mathcal{I}_{ji}] &= \frac{\pi}{4}e^{\frac{i}{2}\pi(\nu-\nu^*)}\mathrm{Re}\bigg[i\int_0^{1}dx\Big(A_i(\epsilon_{\mathrm{IR}}+x)^*+\widetilde{A}_i(\epsilon_{\mathrm{IR}}+x)\Big)\int_0^{\infty}dy\,A_j(\epsilon_{\mathrm{IR}}+ix+iy) \\ &\quad +\int_0^{\infty}dx\Big(A_i(\epsilon_{\mathrm{IR}}+1+ix)^*+\widetilde{A}_i(\epsilon_{\mathrm{IR}}+1+ix)\Big)\int_0^{\infty}dy\,A_j(\epsilon_{\mathrm{IR}}+1+ix+iy)\bigg]+(i\leftrightarrow j)\,.\end{aligned}\tag{A8}$$

By performing similar contour deformations, $\mathrm{Im}[\mathcal{I}_{ij}-\mathcal{I}_{ji}]$ can be also expressed as follows:

$$\begin{aligned}\mathrm{Im}[\mathcal{I}_{ij}-\mathcal{I}_{ji}] &= \frac{\pi}{4}e^{\frac{i}{2}\pi(\nu-\nu^*)}\mathrm{Re}\bigg[\int_0^{1}dx\Big(A_i(\epsilon_{\mathrm{IR}}+x)^*-\widetilde{A}_i(\epsilon_{\mathrm{IR}}+x)\Big)\int_0^{\infty}dy\,A_j(\epsilon_{\mathrm{IR}}+ix+iy) \\ &\quad +\frac{\pi}{4}e^{\frac{i}{2}\pi(\nu-\nu^*)}\mathrm{Im}\bigg[\int_0^{\infty}dx\Big(A_i(\epsilon_{\mathrm{IR}}+1+ix)^*+\widetilde{A}_i(\epsilon_{\mathrm{IR}}+1+ix)\Big)\int_0^{\infty}dy\,A_j(\epsilon_{\mathrm{IR}}+1+ix+iy)\bigg] \\ &\quad -(i\leftrightarrow j)\,.\end{aligned}\tag{A9}$$

The expressions (A8) and (A9) are used in our numerical calculations of the power spectrum.

Appendix B: Analytical calculation of power spectrum for $c_\pi = c_\sigma$

In this appendix we calculate the power spectrum for the case $r_s = c_\sigma/c_\pi = 1$. For this class of models, we can analytically calculate the integrals $\mathcal{I}_{ij}$'s by extending the results in [22]. We first introduce a function $\mathcal{A}(\ell,\nu,x)$ defined by

$$\mathcal{A}(\ell,\nu,x)=x^{-\frac{1}{2}+\ell}e^{ix}H_\nu^{(1)}(x)\,.\tag{B1}$$

In terms of $\mathcal{A}$, A_i can be written as

$$A_1(x)=\mathcal{A}(0,\nu,x)\,,\quad A_2(x)=\mathcal{A}(1,\nu-1,x)\,,\quad A_3(x)=i\mathcal{A}(1,\nu,x)\,,\tag{B2}$$

and hence the x-integrals in the first term and the x_2-integral in the second term of (107) reduce to that of $\mathcal{A}$. Similarly to the case in [22], the indefinite integral $\mathcal{D}$ of $\mathcal{A}$ can be expressed using hyper-geometric functions as

$$\begin{aligned}\mathcal{D}(\ell,\nu\,x) &= \int dx\,\mathcal{A}(\ell,\nu\,x)\\ &= \frac{2^\nu x^{\frac{1}{2}+\ell-\nu}\Gamma(\nu)}{i\pi(\frac{1}{2}+\ell-\nu)}{}_2F_2\Big(\frac{1}{2}-\nu,\frac{1}{2}+\ell-\nu;\frac{3}{2}+\ell-\nu,1-2\nu;+2ix\Big)\\ &\quad+e^{-i\pi\nu}\frac{2^\nu x^{\frac{1}{2}+\ell+\nu}\Gamma(-\nu)}{i\pi(\frac{1}{2}+\ell+\nu)}{}_2F_2\Big(\frac{1}{2}+\nu,\frac{1}{2}+\ell+\nu;\frac{3}{2}+\ell+\nu,1+2\nu;+2ix\Big)\,. \end{aligned}\tag{B3}$$

Then, the integral $\mathcal{I}_{ij}$ can be written as

$$\begin{aligned}\mathcal{I}_{ij} &= \frac{\pi}{8}e^{\frac{i}{2}\pi(\nu-\nu^*)}(-i)^{n_1-\tilde{n}_1+n_2-\tilde{n}_2}\Big(\mathcal{D}(n_1,\nu+n_2,\infty)-\mathcal{D}(n_1,\nu+n_2,0)\Big)^*\Big(\mathcal{D}(\tilde{n}_1,\nu+\tilde{n}_2,\infty)-\mathcal{D}(\tilde{n}_1,\nu+\tilde{n}_2,0)\Big)\\ &\quad-\frac{\pi}{4}i^{\tilde{n}_1+\tilde{n}_2}e^{\frac{i}{2}\pi(\nu-\nu^*)}\int_0^\infty dx_1\widetilde{A}_i(x_1)\Big(\mathcal{D}(\tilde{n}_1,\nu+\tilde{n}_2,\infty)-\mathcal{D}(\tilde{n}_1,\nu+\tilde{n}_2,x_1)\Big)\,,\end{aligned}\tag{B4}$$

where $(n_1,n_2),(\tilde{n}_1,\tilde{n}_2)=(0,0),(1,-1),(1,0)$ for $i,j=1,2,3$, respectively. Here note that $\mathcal{D}(\ell,\nu,x)$ gives a finite value in the limit $x=\infty$ (see appendix B 1 for the derivation):

$$\mathcal{D}(\ell,\nu,\infty) = \frac{1}{i\sqrt{2\pi}}\frac{1}{\sin\pi\nu}\left(\frac{\Gamma(1/2+\ell-\nu)}{\Gamma(1/2-\ell-\nu)}-\frac{\Gamma(1/2+\ell+\nu)}{\Gamma(1/2-\ell+\nu)}\right)\Gamma(-\ell)(-2i)^{-\ell}e^{-\frac{i}{2}\pi(\nu-1/2)}\tag{B5}$$

$$= \frac{\sqrt{\pi}(1-i)e^{-\frac{i}{2}\pi\nu}}{\cos\pi\nu}(-2i)^{-\ell}\frac{1}{\Gamma(\ell+1)}\frac{\Gamma(\frac{1}{2}+\ell+\nu)\Gamma(\frac{1}{2}+\ell-\nu)}{\Gamma(\frac{1}{2}+\nu)\Gamma(\frac{1}{2}-\nu)}\,,\tag{B6}$$

where we used an $i\epsilon$-prescription to drop oscillatory terms $\propto e^{2ix}$ at infinity $x\to\infty$. It should be also noted that the asymptotic behavior of $\mathcal{D}(\ell,\nu,x)$ around $x=0$ is given by

$$\mathcal{D}(\ell,\nu,x) = \frac{2^\nu\Gamma(\nu)}{i\pi(\frac{1}{2}+\ell-\nu)}\Big(x^{\frac{1}{2}+\ell-\nu}+\mathcal{O}(x^{\frac{3}{2}+\ell-\nu})\Big)+e^{-i\pi\nu}\frac{2^\nu\Gamma(-\nu)}{i\pi(\frac{1}{2}+\ell+\nu)}\Big(x^{\frac{1}{2}+\ell+\nu}+\mathcal{O}(x^{\frac{3}{2}+\ell+\nu})\Big)\,,\tag{B7}$$

which is singular for $\mathrm{Re}[1/2+\ell-\nu]<0$. However, as mentioned in appendix A, this kind of singularities cancel out in the calculation of $\mathcal{C}_{ij}$'s and we drop them in the following calculation. We therefore rewrite (B4) as follows:

$$\begin{aligned}\mathcal{I}_{ij} &= \frac{\pi}{8}e^{\frac{i}{2}\pi(\nu-\nu^*)}(-i)^{n_1-\tilde{n}_1+n_2-\tilde{n}_2}\Big(\mathcal{D}(n_1,\nu+n_2,\infty)\Big)^*\mathcal{D}(\tilde{n}_1,\nu+\tilde{n}_2,\infty)\\ &\quad-\frac{\pi}{4}i^{\tilde{n}_1+\tilde{n}_2}e^{\frac{i}{2}\pi(\nu-\nu^*)}\int_0^\infty dx_1\widetilde{A}_i(x_1)\Big(\mathcal{D}(\tilde{n}_1,\nu+\tilde{n}_2,\infty)-\mathcal{D}(\tilde{n}_1,\nu+\tilde{n}_2,x_1)\Big)\\ &= \frac{\pi^2}{4}\frac{1}{\cos^2\pi\nu}2^{-n_1-\tilde{n}_1}(-1)^{n_1-\tilde{n}_1+n_2-\tilde{n}_2}\frac{(\frac{1}{2}+\nu^*+n_2)_{n_1}(\frac{1}{2}-\nu^*-n_2)_{n_1}}{\Gamma(1+n_1)}\frac{(\frac{1}{2}+\nu+\tilde{n}_2)_{\tilde{n}_1}(\frac{1}{2}-\nu-\tilde{n}_2)_{\tilde{n}_1}}{\Gamma(1+\tilde{n}_1)}\\ &\quad-\frac{\pi}{4}i^{\tilde{n}_1+\tilde{n}_2}e^{\frac{i}{2}\pi(\nu-\nu^*)}\int_0^\infty dx_1\widetilde{A}_i(x_1)\Big(\mathcal{D}(\tilde{n}_1,\nu+\tilde{n}_2,\infty)-\mathcal{D}(\tilde{n}_1,\nu+\tilde{n}_2,x_1)\Big)\,,\end{aligned}\tag{B8}$$

with $(a)_m=\dfrac{\Gamma(a+m)}{\Gamma(a)}$.

We next perform the x_1-integral in (B8). To perform this kind of integrals, we use a trick of resummation [22]. Expanding $\widetilde{\mathcal{A}}_i$ in x and using the identity for Bessel functions,

$$x^{-\nu}J_\nu(x)e^{ix}=\sum_{m=0}^\infty a_m^\nu x^m\quad\text{with}\quad a_m^\nu=\frac{2^{m+\nu}i^m\,\Gamma(m+\nu+1/2)}{m!\sqrt{\pi}\,\Gamma(m+2\nu+1)}\,,\tag{B9}$$

$\widetilde{\mathcal{A}}_i$ is rewritten as

$$\begin{aligned}\widetilde{\mathcal{A}}_i &= i^{n_1+n_2}x^{-\frac{1}{2}+n_1}e^{ix}\frac{-i}{\sin\pi(\nu^*+n_2)}\Big(e^{i\pi(\nu^*+n_2)}J_{\nu^*+n_2}(x)-J_{-(\nu^*+n_2)}(x)\Big)\\ &= \frac{-i^{1+n_1+n_2}}{\sin\pi\nu^*}\Big(e^{i\pi\nu^*}\sum_{m=0}^\infty a_m^{\nu^*+n_2}x^{-\frac{1}{2}+m+n_1+n_2+\nu^*}-(-1)^{n_2}\sum_{m=0}^\infty a_m^{-(\nu^*+n_2)}x^{-\frac{1}{2}+m+n_1-n_2-\nu^*}\Big)\,,\end{aligned}\tag{B10}$$

where again $(n_1, n_2) = (0,0), (1,-1), (1,0)$ for $i = 1,2,3$, respectively. Using this expression, the x_1-integral in (B8) can be written as a sum of integrals in the form

$$\int_0^\infty dx\, x^p \left(\mathcal{D}(\ell,\nu,\infty) - \mathcal{D}(\ell,\nu,x)\right) . \tag{B11}$$

Integrating by parts and using $\partial_x \mathcal{D}(\ell,\nu,x) = \mathcal{A}(\ell,\nu,x)$ and $x^{1+p}\mathcal{A}(\ell,\nu,x) = \mathcal{A}(1+\ell+p,\nu,x)$, we rewrite it as

$$\begin{aligned}\int_0^\infty dx\, x^p \left(\mathcal{D}(\ell,\nu,\infty) - \mathcal{D}(\ell,\nu,x)\right) &= \left[\frac{x^{1+p}}{1+p}\left(\mathcal{D}(\ell,\nu,\infty) - \mathcal{D}(\ell,\nu,x)\right)\right]_0^\infty + \int_0^\infty dx\, \frac{1}{1+p}\mathcal{A}(1+\ell+p,\nu,x) \\ &= \frac{1}{1+p}\Big(\mathcal{D}(1+\ell+p,\nu,\infty) + \left[x^{1+p}\mathcal{D}(\ell,\nu,x) - \mathcal{D}(1+\ell+p,\nu,x)\right]_{x\to 0}\Big) \\ &= \frac{1}{1+p}\mathcal{D}(1+\ell+p,\nu,\infty) \,. \end{aligned} \tag{B12}$$

Here we again dropped the contribution from $\mathcal{D}$ at $x = 0$, which vanishes or cancels out in our calculation as mentioned earlier. Then, the x_1-integral can be written as follows:

$$\begin{aligned} &-\frac{\pi}{4} i^{\tilde{n}_1+\tilde{n}_2} e^{\frac{i}{2}\pi(\nu-\nu^*)} \int_0^\infty dx_1 \widetilde{A}_i(x_1) \Big(\mathcal{D}(\tilde{n}_1, \nu+\tilde{n}_2, \infty) - \mathcal{D}(\tilde{n}_1, \nu+\tilde{n}_2, x_1)\Big) \\ &= \frac{\pi}{4}\frac{i^{1+n_1+\tilde{n}_1+\tilde{n}_2}}{\sin \pi\nu^*}\Bigg(e^{\frac{i}{2}\pi(\nu+\nu^*+n_2)} \sum_{m=0}^{\infty} \frac{a_m^{\nu^*+n_2}}{\frac{1}{2}+m+n_1+n_2+\nu^*}\mathcal{D}(1/2+m+n_1+n_2+\nu^*+\tilde{n}_1, \nu+\tilde{n}_2, \infty) \\ &\qquad\qquad - e^{\frac{i}{2}\pi(\nu-\nu^*-n_2)} \sum_{m=0}^{\infty} \frac{a_m^{-(\nu^*+n_2)}}{\frac{1}{2}+m+n_1-n_2-\nu^*}\mathcal{D}(1/2+m+n_1-n_2-\nu^*+\tilde{n}_1, \nu+\tilde{n}_2, \infty)\Bigg) \,. \end{aligned} \tag{B13}$$

After some lengthy calculations, we obtain

$$\begin{aligned} &\frac{\pi}{4}\frac{i^{1+n_1+\tilde{n}_1+\tilde{n}_2}}{\sin \pi\nu^*} e^{\frac{i}{2}\pi(\nu\pm\nu^*\pm n_2)} \frac{a_m^{\pm(\nu^*+n_2)}}{\frac{1}{2}+m+n_1\pm n_2\pm\nu^*}\mathcal{D}(1/2+m+n_1\pm n_2\pm\nu^*+\tilde{n}_1, \nu+\tilde{n}_2, \infty) \\ &= \frac{i\, e^{\pm i\pi\nu^*}}{\sin \pi\nu^*} \frac{2^{-n_1-\tilde{n}_1-2}(-1)^{m+\tilde{n}_2}}{\left(\frac{1}{2}+m+n_1\pm(\nu^*+n_2)\right)\left(\frac{1}{2}+m\pm(\nu^*+n_2)\right)_{1+n_1+\tilde{n}_1}} \\ &\quad \times \begin{cases} (m+1)_{n_1\pm n_2+\tilde{n}_1\mp\tilde{n}_2} \left(-m-n_1\mp n_2-\tilde{n}_1\mp\tilde{n}_2\mp 2\nu^*\right)_{n_1\mp n_2+\tilde{n}_1\pm\tilde{n}_2} & \text{for real } \nu \,, \\ (m+1)_{n_1\pm n_2+\tilde{n}_1\pm\tilde{n}_2} \left(-m-n_1\mp n_2-\tilde{n}_1\pm\tilde{n}_2\mp 2\nu^*\right)_{n_1\mp n_2+\tilde{n}_1\mp\tilde{n}_2} & \text{for pure-imaginary } \nu \,, \end{cases} \end{aligned} \tag{B14}$$

where $\Gamma(z)\Gamma(1-z) = \dfrac{\pi}{\sin \pi z}$ was used. Finally, it is necessary to re-sum (B14) with respect to m. In the case of $\mathrm{Re}[\mathcal{I}_{11}]$ ($n_i = \tilde{n}_i = 0$), for example, we perform the resummation as follows:

$$\begin{aligned} &-\frac{\pi}{4} e^{\frac{i}{2}\pi(\nu-\nu^*)} \int_0^\infty dx_1 \widetilde{A}_1(x_1) \Big(\mathcal{D}(0,\nu,\infty) - \mathcal{D}(0,\nu,x_1)\Big) \\ &= \frac{i}{4\sin \pi\nu} \sum_{m=0}^{\infty} \left[\frac{(-1)^m e^{i\pi\nu}}{(m+\frac{1}{2}+\nu)^2} - \frac{(-1)^m e^{-i\pi\nu}}{(m+\frac{1}{2}-\nu)^2}\right] \\ &= \frac{i}{4\sin \pi\nu}\left[e^{i\pi\nu}\Phi(-1,2,\tfrac{1}{2}+\nu) - e^{-i\pi\nu}\Phi(-1,2,\tfrac{1}{2}-\nu)\right] \\ &= -\frac{i\, e^{i\pi\nu}}{16\sin \pi\nu}\left[\psi^{(1)}\Big(\frac{3}{4}+\frac{\nu}{2}\Big) - \psi^{(1)}\Big(\frac{1}{4}+\frac{\nu}{2}\Big)\right] + \frac{i\, e^{-i\pi\nu}}{16\sin \pi\nu}\left[\psi^{(1)}\Big(\frac{3}{4}-\frac{\nu}{2}\Big) - \psi^{(1)}\Big(\frac{1}{4}-\frac{\nu}{2}\Big)\right] . \end{aligned}$$

Here $\Phi(z,s,\alpha)$ and $\psi^{(n)}(z)$ are the Lerch transcendent and the polygamma function, respectively:

$$\Phi(z,s,\alpha) = \sum_{m=0}^{\infty} \frac{z^m}{(m+\alpha)^s} \,, \quad \psi^{(n)}(z) = (-1)^{n+1} n! \sum_{m=0}^{\infty} \frac{1}{(m+z)^{n+1}} \,, \tag{B15}$$

which satisfy the following relation:

$$\Phi(-1, n+1, z) = (-1)^n \, n! \, 2^{-n-1} \left[\psi^{(n)}\Big(\frac{z+1}{2}\Big) - \psi^{(n)}\Big(\frac{z}{2}\Big) \right] . \tag{B16}$$

We then obtain

$$\mathrm{Re}[\mathcal{I}_{11}] = \frac{\pi^2}{4\cos^2 \pi\nu} + \mathrm{Re}\left[\frac{i\, e^{-i\pi\nu}}{16 \sin \pi\nu} \Big[\psi^{(1)}\Big(\frac{3}{4} + \frac{\nu}{2}\Big) - \psi^{(1)}\Big(\frac{1}{4} + \frac{\nu}{2}\Big) \Big] - \frac{i\, e^{i\pi\nu}}{16 \sin \pi\nu} \Big[\psi^{(1)}\Big(\frac{3}{4} - \frac{\nu}{2}\Big) - \psi^{(1)}\Big(\frac{1}{4} - \frac{\nu}{2}\Big) \Big] \right] , \tag{B17}$$

which reproduces the result in [22]. In a similar way, we can obtain analytic expression for $\mathcal{I}_{ij}$'s, and the results are summarized as follows:

$$\mathcal{I}_{22} = \begin{cases} \dfrac{\pi^2 |\nu - \frac{1}{2}|^2 |\nu - \frac{3}{2}|^2}{16 \cos^2 \pi\nu} + \dfrac{1}{16} - \dfrac{i}{128 \sin \pi\nu} \Big(e^{i\pi\nu} (2\nu - 3)(2\nu - 5) - e^{-i\pi\nu} (2\nu + 1)(2\nu - 1) \Big) \\ + \dfrac{i}{256 \sin \pi\nu} (2\nu - 1)^2 (2\nu - 3)^2 \Big(e^{i\pi\nu} \Phi(-1, 2, \tfrac{1}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 2, \tfrac{5}{2} - \nu) \Big) \quad \text{for real } \nu , \\ \\ \dfrac{\pi^2 |\nu - \frac{1}{2}|^2 |\nu - \frac{3}{2}|^2}{16 \cos^2 \pi\nu} + \dfrac{1}{16} - \dfrac{i}{128 \sin \pi\nu} \Big(e^{i\pi\nu} \dfrac{-3 - 82\nu + 12\nu^2 + 72\nu^3}{3 + 2\nu} + e^{-i\pi\nu} \dfrac{-15 - 126\nu - 36\nu^2 + 56\nu^3}{1 + 2\nu} \Big) \\ + \dfrac{i}{8 \sin \pi\nu} \nu(-3 + 4\nu^2) \Big(e^{i\pi\nu} \, \Phi(-1, 1, \tfrac{3}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 1, -\tfrac{1}{2} - \nu) \Big) \\ + \dfrac{i}{256 \sin \pi\nu} (4\nu^2 - 1)(4\nu^2 - 9) \Big(e^{i\pi\nu} \Phi(-1, 2, \tfrac{5}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 2, \tfrac{1}{2} - \nu) \Big) \quad \text{for pure-imaginary } \nu , \end{cases} \tag{B18}$$

$$\begin{aligned} \mathcal{I}_{33} = {} & \frac{\pi^2 |\nu + \frac{1}{2}|^2 |\nu - \frac{1}{2}|^2}{16 \cos^2 \pi\nu} - \frac{1}{16} + \frac{i}{128 \sin \pi\nu} \Big(e^{i\pi\nu} (2\nu - 1)(2\nu - 3) - e^{-i\pi\nu} (2\nu + 1)(2\nu + 3) \Big) \\ & - \frac{i}{256 \sin \pi\nu} (2\nu + 1)^2 (2\nu - 1)^2 \Big(e^{i\pi\nu} \Phi(-1, 2, \tfrac{3}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 2, \tfrac{3}{2} - \nu) \Big) , \end{aligned} \tag{B19}$$

$$\begin{aligned} \mathcal{I}_{12} = {} & -\frac{\pi^2 (\nu - \frac{1}{2})(\nu - \frac{3}{2})}{8 \cos^2 \pi\nu} - \frac{i}{16 \sin \pi\nu} \Big(e^{i\pi\nu} (2\nu - 1) + e^{-i\pi\nu} (2\nu + 1) \Big) \\ & + \frac{i}{16 \sin \pi\nu} (-3 + 4\nu^2) \Big(e^{i\pi\nu} \Phi(-1, 1, \tfrac{1}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 1, \tfrac{1}{2} - \nu) \Big) \\ & - \frac{i}{32 \sin \pi\nu} (-3 + 2\nu)(-1 + 2\nu) \Big(e^{i\pi\nu} \Phi(-1, 2, \tfrac{1}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 2, \tfrac{1}{2} - \nu) \Big) , \end{aligned} \tag{B20}$$

$$\begin{aligned} \mathcal{I}_{21} = {} & -\frac{\pi^2 (\nu^* - \frac{1}{2})(\nu^* - \frac{3}{2})}{8 \cos^2 \pi\nu} + \frac{i(-1 + 4\nu^2)}{32 \sin \pi\nu^*} \Big(e^{i\pi\nu^*} \Phi(-1, 1, -\tfrac{1}{2} + \nu^*) - e^{-i\pi\nu^*} \Phi(-1, 1, \tfrac{3}{2} - \nu^*) \Big) \\ & - \frac{i(4\nu^2 - 5)}{32 \sin \pi\nu^*} \Big(e^{i\pi\nu^*} \Phi(-1, 1, \tfrac{1}{2} + \nu^*) - e^{-i\pi\nu^*} \Phi(-1, 1, \tfrac{5}{2} - \nu^*) \Big) \\ & - \frac{i(-3 + 2\nu^*)(-1 + 2\nu^*)}{32 \sin \pi\nu^*} \Big(e^{i\pi\nu^*} \Phi(-1, 2, \tfrac{1}{2} + \nu^*) - e^{-i\pi\nu^*} \Phi(-1, 2, \tfrac{5}{2} - \nu^*) \Big) , \end{aligned} \tag{B21}$$

$$\begin{aligned} \mathcal{I}_{13} = {} & \frac{\pi^2 (\nu + \frac{1}{2})(\nu - \frac{1}{2})}{8 \cos^2 \pi\nu} + \frac{i}{16 \sin \pi\nu} \Big(e^{i\pi\nu} (2\nu - 1) + e^{-i\pi\nu} (2\nu + 1) \Big) \\ & - \frac{i(1 + 4\nu^2)}{16 \sin \pi\nu} \Big(e^{i\pi\nu} \Phi(-1, 1, \tfrac{1}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 1, \tfrac{1}{2} - \nu) \Big) \\ & + \frac{i(-1 + 4\nu^2)}{32 \sin \pi\nu} \Big(e^{i\pi\nu} \Phi(-1, 2, \tfrac{1}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 2, \tfrac{1}{2} - \nu) \Big) , \end{aligned} \tag{B22}$$

$$\begin{aligned} \mathcal{I}_{31} = {} & \frac{\pi^2 (\nu^* + \frac{1}{2})(\nu^* - \frac{1}{2})}{8 \cos^2 \pi\nu} + \frac{i(-1 + 4\nu^2)}{32 \sin \pi\nu} \Big(e^{i\pi\nu} \Phi(-1, 1, \tfrac{1}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 1, \tfrac{1}{2} - \nu) \Big) \\ & - \frac{i(3 + 4\nu^2)}{32 \sin \pi\nu} \Big(e^{i\pi\nu} \Phi(-1, 1, \tfrac{3}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 1, \tfrac{3}{2} - \nu) \Big) \\ & - \frac{i(-1 + 4\nu^2)}{32 \sin \pi\nu} \Big(e^{i\pi\nu} \Phi(-1, 2, \tfrac{3}{2} + \nu) - e^{-i\pi\nu} \Phi(-1, 2, \tfrac{3}{2} - \nu) \Big) , \end{aligned} \tag{B23}$$

$$\begin{aligned}\mathcal{I}_{23} = &-\frac{\pi^2|\nu-\frac{1}{2}|^2(\nu+\frac{1}{2})(\nu^*-\frac{3}{2})}{16\cos^2\pi\nu}+\frac{1}{16}-\frac{i}{128\sin\pi\nu^*}\Big(e^{i\pi\nu^*}(-11+12\nu^2)+e^{-i\pi\nu^*}\frac{-1+2\nu^*}{-3+2\nu^*}(1-20\nu^*+20\nu^2)\Big)\,,\\ &+\frac{i(-1+2\nu^*)(-1-4\nu^*+4\nu^2)}{32\sin\pi\nu^*}\Big(e^{i\pi\nu^*}\Phi(-1,1,-\tfrac{1}{2}+\nu^*)-e^{-i\pi\nu^*}\Phi(-1,1,\tfrac{3}{2}-\nu^*)\Big)\\ &-\frac{i(-1+2\nu^*)^2(-3+2\nu^*)(1+2\nu^*)}{256\sin\pi\nu^*}\Big(e^{i\pi\nu^*}\Phi(-1,2,\tfrac{1}{2}+\nu^*)-e^{-i\pi\nu^*}\Phi(-1,2,\tfrac{5}{2}-\nu^*)\Big)\,,\end{aligned} \tag{B24}$$

$$\begin{aligned}\mathcal{I}_{32} = &-\frac{\pi^2|\nu-\frac{1}{2}|^2(\nu-\frac{3}{2})(\nu^*+\frac{1}{2})}{16\cos^2\pi\nu}-\frac{1}{16}-\frac{i}{128\sin\pi\nu}\Big(e^{i\pi\nu}\frac{-1+2\nu}{1+2\nu}(1-20\nu+20\nu^2)+e^{-i\pi\nu}(1-24\nu+12\nu^2)\Big)\\ &+\frac{i(-1+2\nu)(-1-4\nu+4\nu^2)}{32\sin\pi\nu}\Big(e^{i\pi\nu}\Phi(-1,1,\tfrac{1}{2}+\nu)-e^{-i\pi\nu}\Phi(-1,1,\tfrac{1}{2}-\nu)\Big)\\ &+\frac{i(-3+2\nu)(-1+2\nu)^2(1+2\nu)}{256\sin\pi\nu}\Big(e^{i\pi\nu}\Phi(-1,2,\tfrac{3}{2}+\nu)-e^{-i\pi\nu}\Phi(-1,2,\tfrac{3}{2}-\nu)\Big)\,.\end{aligned} \tag{B25}$$

Here we used $\sum_{m=0}^{\infty}(-1)^m = \frac{1}{1+1} = \frac{1}{2}$. As we displayed in figure 1, the analytic results obtained in this appendix and the numerical results for $r_s = 1$ well coincide with each other.

1. Asymptotic behavior of $\mathcal{D}(\ell,\nu,x)$

In this subsection we derive the asymptotic behavior (B5) in the limit $x \to \infty$ of the function $\mathcal{D}(\ell,\nu,x)$:

$$\begin{aligned}\mathcal{D}(\ell,\nu\, x) = &\frac{2^\nu x^{\frac{1}{2}+\ell-\nu}\Gamma(\nu)}{i\pi(\frac{1}{2}+\ell-\nu)}{}_2F_2\Big(\frac{1}{2}-\nu,\frac{1}{2}+\ell-\nu;\frac{3}{2}+\ell-\nu,1-2\nu;2ix\Big)\\ &+e^{-i\pi\nu}\frac{2^{-\nu}x^{\frac{1}{2}+\ell+\nu}\Gamma(-\nu)}{i\pi(\frac{1}{2}+\ell+\nu)}{}_2F_2\Big(\frac{1}{2}+\nu,\frac{1}{2}+\ell+\nu;\frac{3}{2}+\ell+\nu,1+2\nu;2ix\Big)\,.\end{aligned} \tag{B26}$$

We use the following asymptotic expansion of hypergeometric functions:

$$\begin{aligned}{}_2F_2(a_1,a_2;b_1,b_2;z) = &\frac{\Gamma(b_1)\Gamma(b_2)}{\Gamma(a_1)\Gamma(a_2)}e^z z^{a_1+a_2-b_1-b_2}\sum_{k=0}^{\infty}c_k z^{-k}\\ &+\frac{\Gamma(b_1)\Gamma(b_2)\Gamma(a_2-a_1)}{\Gamma(a_2)\Gamma(b_1-a_1)\Gamma(b_2-a_1)}(-z)^{-a_1}{}_3F_1(a_1,a_1-b_1+1,a_1-b_2+1;a_1-a_2+1;-1/z)\\ &+\frac{\Gamma(b_1)\Gamma(b_2)\Gamma(a_1-a_2)}{\Gamma(a_1)\Gamma(b_1-a_2)\Gamma(b_2-a_2)}(-z)^{-a_2}{}_3F_1(a_2,a_2-b_1+1,a_2-b_2+1;a_2-a_1+1;-1/z)\,,\end{aligned} \tag{B27}$$

where c_k's are numerical factors independent of z. The first term gives an oscillating term $\sim e^{2ix}$, which can be dropped by an $i\epsilon$-prescription. Then, let us consider the contribution of the last two terms. They are respectively in the form

$$\begin{aligned}&\frac{1}{i\sqrt{2\pi}\ell}\frac{1}{\sin\pi\nu}x^\ell e^{-\frac{i}{2}\pi(\nu-1/2)}{}_3F_1(1/2+\nu,1/2-\nu,-\ell;1-\ell;i/(2x))\\ &+\frac{1}{i\sqrt{2\pi}}\frac{1}{\sin\pi\nu}\frac{\Gamma(1/2+\ell-\nu)}{\Gamma(1/2-\ell-\nu)}\Gamma(-\ell)(-2i)^{-\ell}e^{-\frac{i}{2}\pi(\nu-1/2)}{}_3F_1(1/2+\ell+\nu,1/2+\ell-\nu,0;1+\ell;i/(2x))\,,\end{aligned} \tag{B28}$$

and

$$\begin{aligned}&-\frac{1}{i\sqrt{2\pi}\ell}\frac{1}{\sin\pi\nu}x^\ell e^{-\frac{i}{2}\pi(\nu-1/2)}{}_3F_1(1/2+\nu,1/2-\nu,-\ell;1-\ell;i/(2x))\\ &-\frac{1}{i\sqrt{2\pi}}\frac{1}{\sin\pi\nu}\frac{\Gamma(1/2+\ell+\nu)}{\Gamma(1/2-\ell+\nu)}\Gamma(-\ell)(-2i)^{-\ell}e^{-\frac{i}{2}\pi(\nu-1/2)}{}_3F_1(1/2+\ell+\nu,1/2+\ell-\nu,0;1+\ell;i/(2x))\,,\end{aligned} \tag{B29}$$

where note that hypergeometric functions ${}_pF_q(a_1,\dots,a_p;b_1,\dots,b_q;z)$ are symmetric under the permutations of a_i's and those of b_i's, respectively. The first term in (B28) and that in (B29) cancel out and we obtain

$$\begin{aligned}\mathcal{D}(\ell,\nu,x) =& \frac{1}{i\sqrt{2\pi}}\frac{1}{\sin\pi\nu}\frac{\Gamma(1/2+\ell-\nu)}{\Gamma(1/2-\ell-\nu)}\Gamma(-\ell)(-2i)^{-\ell}e^{-\frac{i}{2}\pi(\nu-1/2)}{}_3F_1(1/2+\ell+\nu,1/2+\ell-\nu,0;1+\ell;i/(2x))\\ &-\frac{1}{i\sqrt{2\pi}}\frac{1}{\sin\pi\nu}\frac{\Gamma(1/2+\ell+\nu)}{\Gamma(1/2-\ell+\nu)}\Gamma(-\ell)(-2i)^{-\ell}e^{-\frac{i}{2}\pi(\nu-1/2)}{}_3F_1(1/2+\ell+\nu,1/2+\ell-\nu,0;1+\ell;i/(2x))\,,\end{aligned} \tag{B30}$$

where note that we did not use any approximation so far. Finally, taking the limit $x\to\infty$, we conclude that

$$\mathcal{D}(\ell,\nu,\infty) = \frac{1}{i\sqrt{2\pi}}\frac{1}{\sin\pi\nu}\left(\frac{\Gamma(1/2+\ell-\nu)}{\Gamma(1/2-\ell-\nu)}-\frac{\Gamma(1/2+\ell+\nu)}{\Gamma(1/2-\ell+\nu)}\right)\Gamma(-\ell)(-2i)^{-\ell}e^{-\frac{i}{2}\pi(\nu-1/2)}\,, \tag{B31}$$

where we used ${}_3F_1(a_1,a_2,a_3;b;0)=1$. Using the identity

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin\pi z}\,, \tag{B32}$$

we can also rewrite (B31) as follows:

$$\mathcal{D}(\ell,\nu,\infty) = \frac{\sqrt{\pi}(1-i)e^{-\frac{i}{2}\pi\nu}}{\cos\pi\nu}(-2i)^{-\ell}\frac{1}{\Gamma(\ell+1)}\frac{\Gamma(\frac{1}{2}+\ell+\nu)\Gamma(\frac{1}{2}+\ell-\nu)}{\Gamma(\frac{1}{2}+\nu)\Gamma(\frac{1}{2}-\nu)}\,, \tag{B33}$$

which reproduces the result in [22] for $\ell=0$.

[1] K. Sato, Mon. Not. Roy. Astron. Soc. **195**, 467 (1981). A. H. Guth, Phys. Rev. D **23**, 347 (1981);
[2] A. A. Starobinsky, Phys. Lett. B **91**, 99 (1980);
[3] E. Komatsu *et al.* [WMAP Collaboration],
[4] T. Kobayashi, M. Yamaguchi and J. Yokoyama, Prog. Theor. Phys. **126**, 511 (2011) [arXiv:1105.5723 [hep-th]].
[5] G. W. Horndeski, Int. J. Theor. Phys. 10 (1974) 363-384.
[6] C. Charmousis, E. J. Copeland, A. Padilla and P. M. Saffin, Phys. Rev. Lett. **108**, 051101 (2012) [arXiv:1106.2000 [hep-th]].
[7] C. Deffayet, X. Gao, D. A. Steer and G. Zahariade, Phys. Rev. D **84**, 064039 (2011) [arXiv:1103.3260 [hep-th]].
[8] T. Kobayashi, M. Yamaguchi and J. Yokoyama, Phys. Rev. Lett. **105**, 231302 (2010) [arXiv:1008.0603 [hep-th]]; C. Burrage, C. de Rham, D. Seery and A. J. Tolley, JCAP **1101**, 014 (2011) [arXiv:1009.2497 [hep-th]].
[9] S. Mizuno, K. Koyama, Phys. Rev. **D82**, 103518 (2010). [arXiv:1009.0677 [hep-th]]; A. De Felice and S. Tsujikawa, JCAP **1104**, 029 (2011) [arXiv:1103.1172 [astro-ph.CO]]; T. Kobayashi, M. Yamaguchi and J. Yokoyama, Phys. Rev. D **83**, 103524 (2011) [arXiv:1103.1740 [hep-th]].
[10] X. Gao and D. A. Steer, JCAP **1112**, 019 (2011) [arXiv:1107.2642 [astro-ph.CO]]; A. De Felice and S. Tsujikawa, Phys. Rev. D **84**, 083504 (2011) [arXiv:1107.3917 [gr-qc]]; S. Renaux-Petel, JCAP **1202**, 020 (2012) [arXiv:1107.5020 [astro-ph.CO]].
[11] X. Gao, T. Kobayashi, M. Yamaguchi and J. 'i. Yokoyama, Phys. Rev. Lett. **107**, 211301 (2011) [arXiv:1108.3513 [astro-ph.CO]]; X. Gao, T. Kobayashi, M. Shiraishi, M. Yamaguchi, J. 'i. Yokoyama and S. Yokoyama, arXiv:1207.0588 [astro-ph.CO].
[12] M. Yamaguchi and J. 'i. Yokoyama, Phys. Rev. D **74**, 043523 (2006) [hep-ph/0512318].
[13] S. Cremonini, Z. Lalak and K. Turzynski, JCAP **1103**, 016 (2011) [arXiv:1010.3021 [hep-th]]. A. Avgoustidis, S. Cremonini, A. -C. Davis, R. H. Ribeiro, K. Turzynski and S. Watson, JCAP **1206**, 025 (2012) [arXiv:1203.0016 [hep-th]].
[14] A. Achucarro, J. -O. Gong, S. Hardeman, G. A. Palma and S. P. Patil, JCAP **1101**, 030 (2011) [arXiv:1010.3693 [hep-ph]].
[15] G. Shiu and J. Xu, Phys. Rev. D **84**, 103509 (2011) [arXiv:1108.0981 [hep-th]].
[16] S. Cespedes, V. Atal and G. A. Palma, JCAP **1205**, 008 (2012) [arXiv:1201.4848 [hep-th]].
[17] A. Achucarro, J. -O. Gong, S. Hardeman, G. A. Palma and S. P. Patil, JHEP **1205**, 066 (2012) [arXiv:1201.6342 [hep-th]].
[18] S. Pi and M. Sasaki, JCAP **1210**, 051 (2012) [arXiv:1205.0161 [hep-th]].
[19] X. Gao, D. Langlois and S. Mizuno, arXiv:1205.5275 [hep-th].
[20] R. Saito, M. Nakashima, Y. -i. Takamizu and J. 'i. Yokoyama, arXiv:1206.2164 [astro-ph.CO].
[21] X. Chen and Y. Wang, Phys. Rev. D **81**, 063511 (2010) [arXiv:0909.0496 [astro-ph.CO]]; X. Chen and Y. Wang, JCAP **1004**, 027 (2010) [arXiv:0911.3380 [hep-th]];
[22] X. Chen and Y. Wang, arXiv:1205.0160 [hep-th].
[23] E. J. Copeland, A. R. Liddle, D. H. Lyth, E. D. Stewart and D. Wands, Phys. Rev. D **49**, 6410 (1994) [astro-ph/9401011]; E. D. Stewart, Phys. Rev. D **51**, 6847 (1995) [hep-ph/9405389].
[24] D. Baumann and D. Green, Phys. Rev. D **85**, 103520 (2012) [arXiv:1109.0292 [hep-th]].

[25] L. McAllister, S. Renaux-Petel and G. Xu, JCAP **1210**, 046 (2012) [arXiv:1207.0317 [astro-ph.CO]].

[26] A. Mazumdar and J. Rocher, Phys. Rept. **497**, 85 (2011) [arXiv:1001.0993 [hep-ph]]; M. Yamaguchi, Class. Quant. Grav. **28**, 103001 (2011) [arXiv:1101.2488 [astro-ph.CO]].

[27] C. Cheung, P. Creminelli, A. L. Fitzpatrick, J. Kaplan and L. Senatore, JHEP **0803**, 014 (2008) [arXiv:0709.0293 [hep-th]];

[28] L. Senatore and M. Zaldarriaga, JHEP **1204**, 024 (2012) [arXiv:1009.2093 [hep-th]].

[29] S. Weinberg, Phys. Rev. D **77**, 123541 (2008) [arXiv:0804.4291 [hep-th]].

[30] J. M. Maldacena, JHEP **0305**, 013 (2003) [astro-ph/0210603].

[31] S. Groot Nibbelink and B. J. W. van Tent, Class. Quant. Grav. **19**, 613 (2002) [hep-ph/0107272].

[32] E. Sefusatti, J. R. Fergusson, X. Chen and E. P. S. Shellard, JCAP **1208**, 033 (2012) [arXiv:1204.6318 [astro-ph.CO]].